Petroleum Resource Assessment of the Yukon North Coast, Yukon Territory, Canada

by
P.K. Hannigan

March, 2001
Whitehorse, Yukon

Yukon
Economic Development

© March, 2001
Oil and Gas Resources Branch
Department of Economic Development
Government of the Yukon
Box 2703
Whitehorse, Yukon Y1A 2C6
phone: 867-667-3427
fax: 867-393-6262
website: www.yukonoilandgas.com

ISBN 1-55018-997-2

FOREWORD

On November 19, 1998, the Government of Canada transferred to the Government of Yukon the administrative legislative powers and responsibilities of managing onshore oil and gas resources. Yukon oil and gas resources are now governed under the Yukon *Oil and Gas Act*.

A study of the petroleum resources of the Old Crow Basin in the Yukon Territory was undertaken by Geological Survey of Canada (GSC) in response to a request from the Government of Yukon. Assessment of petroleum resource potential is important for forming regulatory policies for these resources and for providing a basis for planning and issuing exploration rights.

102-05696/

EXECUTIVE SUMMARY

This study was undertaken by the Geological Survey of Canada on behalf of the Yukon government as part of its ongoing oil and gas resources management program. The objective of the study was to investigate the hydrocarbon resource potential of the northern Arctic Coast in the Yukon Territory. A quantitative analysis was utilized to derive a numerical estimate of resources that may exist in the area. Due to the scarcity of defined pools with established reserves, probability distributions of reservoir parameters and marginal risk factors were employed to generate a range of hydrocarbon potential estimates. The predicted wide ranges of resource potential reveal the uncertainties involved in the analysis of frontier exploration plays.

Numerous regional tectonic and structural elements occur beneath the coastal lowland and shallow waters of the Beaufort Sea on the North Coast of the Yukon. The area west of the northern Richardson Mountains and extending into the Beaufort Sea consists of highly deformed strata of the Cordilleran Foldbelt. Structures, including compressional folds and strike-slip faults formed during the Laramide Orogeny, are underlain by older tectonic features such as the Cache Creek Uplift and Rapid Depression which originated as Mesozoic fault-bounded structures. North of Rapid Depression, several significant large-scale tectonic features, such as Blow River and Herschel highs, are cored by closely spaced anticlines. On the south flank of Herschel High, there lies a large synclinal structure filled with Eocene and younger strata called the Demarcation Subbasin.

The hydrocarbon potential volumes were derived using the Geological Survey of Canada's PETRIMES assessment methodology system. This resource study embraced analyses of si : conceptual and immature plays, each of which incorporated estimations of field size parameters, numbers of prospects and exploration risks. The median estimate for total oil and gas potential for all coastal lowland plays are 39.8 million m^3 and 38.6 billion m^3, respectively. There are no discovered reserves in the Yukon portion of the assessment region. The assessment results indicate that four gas fields greater than 3,000 million m^3 (100 BCF) are expected in the region. No oil fields larger than 160 million m^3 (1 billion bbls.) are predicted beneath Yukon North Coast. Even though geological risk factors are substantial, significant gas potential is predicted for the Herschel play.

CONTENTS

Introduction **1**
Acknowledgements 1
Terminology 1
Method and content 2

Resource assessment procedure **3**
Geological play definition 3
Compilation of play data 3
Subjective probability analysis 3

Regional geology **5**
Stratigraphy and depositional setting 7
Inuvikian (Proterozoic) 7
Franklinian (Lower Paleozoic) 7
Ellesmerian (Carboniferous to middle Hauterivian) 9
Brookian (Upper Hauterivian to Eocene) 11
Structural geology 12

Petroleum geology **13**
Exploration history 13
Reservoirs 13
Carboniferous 13
Permo-Triassic 13
Jurassic 14
Cretaceous 14
Tertiary 14
Seals 15
Traps 15
Source rocks 15
Timing of hydrocarbon generation 16
Hydrocarbon shows 17

Hydrocarbon assessment **18**
Hydrocarbon plays 19
South Delta–Mesozoic oil and gas play 19
South Delta–Paleozoic gas play 27
Herschel oil and gas play 30
Yukon coastal plain gas play 36
Discussion of assessment results 40
Resource potential 40
Resource distributions 41
Assessment results and exploration history 43

Conclusions **44**

References **45**

Appendix 1: Input data for North Coast hydrocarbon assessments **50**

Appendix 2: Output for North Coast hydrocarbon assessments **57**

INTRODUCTION

This study was undertaken by the Geological Survey of Canada on behalf of the Yukon Territorial Government as part of its ongoing oil and gas resource management program. The objective of this study was to investigate the hydrocarbon resource potential of the North Coast in the Yukon Territory (Fig. 1). A quantitative analysis was utilized to derive a numerical estimate of resources that may exist in the region. Due to the scarcity of defined pools with established reserves, probability distributions of reservoir parameters and marginal risk factors were adopted to generate a range of hydrocarbon potential estimates indicative of the uncertainties involved in the analysis of frontier conceptual and immature plays.



Figure 1. *Map of northwestern Canada showing study area location.*

Regional petroleum resource assessments have been prepared periodically for numerous sedimentary basins in Canada by the Geological Survey of Canada. These studies incorporate systematic basin analysis with subsequent statistical resource evaluations (Podruski, *et al.*, 1988; Wade, *et al.*, 1989; Sinclair, *et al.*, 1992; Reinson, *et al.*, 1993; Bird, *et al.*, 1994; Dixon, *et al.*, 1994; Hannigan, *et al.*, 1998, 1999; Hannigan, in press (a), (b) and (c)).

This report provides an overview of the petroleum geology of the Yukon North Coast region and presents quantitative estimates of the oil and gas resources contained therein. The geological and resource framework for the region will assist government agencies in evaluating land-use and moratorium issues and petroleum industry companies in pursuing future exploration opportunities.

ACKNOWLEDGEMENTS

The author would like to acknowledge staff at GSC-Calgary for their insight and geological expertise on this area, specifically, J. Dixon, J.R. Dietrich and K.G. Osadetz.

TERMINOLOGY

The terminology and procedures used in this report follow those outlined in Reinson, *et al.* (1993) and are summarized below.

Oil is defined as any naturally occurring liquid that, at the conditions under which it is measured or estimated, is primarily composed of hydrocarbon molecules and is readily producible from a borehole.

Natural gas is defined as any gas (at standard pressure and temperature, 101.33 kPa and 15oC) of natural origin, comprised mostly of hydrocarbon molecules, producible from a borehole (Potential Gas Committee, 1990). Natural gas may contain significant amounts of non-hydrocarbon gas such as H2S, CO2 or He. In this study, non-hydrocarbon gas was not considered due to lack of information on gas compositions in these basins.

Condensate is a mixture, mainly of pentanes and heavier hydrocarbons, that can be recovered from an underground reservoir that may be gaseous in its virgin reservoir state

but is liquid at the conditions under which its volume is measured (Alberta Energy and Utilities Board, 1999).

Raw gas is unprocessed natural gas, containing methane, inert and acid gases, impurities and other hydrocarbons, some of which can be recovered as liquids. *Non-associated gas* is natural gas that is not in contact with oil in a reservoir. *Associated gas* is natural gas that occurs in oil reservoirs as free gas. *Solution gas* is natural gas that is dissolved in crude oil in reservoirs. In this report, insufficient information is available in order to differentiate non-associated, associated, and solution gas. All gas figures reported represent initial raw gas volumes.

Resource indicates all hydrocarbon accumulations known or inferred to exist. *Resource, resource endowment and endowment* are synonymous and can be used interchangeably. *Reserves* are that portion of the resource that has been discovered, while *potential* represents the portion of the resource that is not discovered but is inferred to exist. The terms *potential* and *undiscovered resources* are synonymous and may be used interchangeably.

Gas-in-place indicates the gas volume found in the ground, regardless of what portion is recoverable. *Initial in-place volume* is the gross volume of raw gas, before production. *Recoverable in-place volume* represents the volume expected to be recovered using current technology and costs. These definitions can be applied to oil volumes as well.

A *prospect* is defined as an untested exploration target within a single stratigraphic interval; it may or may not contain hydrocarbons. A prospect is not synonymous with an undiscovered pool. An undiscovered pool is a prospect that contains hydrocarbons but has not been tested as yet. A *pool* is defined as a discovered accumulation of oil or gas typically within a single stratigraphic interval that is separate, hydrodynamically or otherwise, from another hydrocarbon accumulation. A *field* consists of one or more oil and/or gas pools within a single structure or trap. Similar to most frontier regions, the assessment of petroleum resources of the Yukon North Coast is based on estimates of field rather than pool sizes. A *play* is defined as a family of pools and/or prospects that share a common history of hydrocarbon generation, migration, reservoir development and trap configuration.

Plays are grouped into two categories: *established* and *conceptual* plays. *Established plays* are demonstrated to exist due to the discovery of pools with established reserves. *Conceptual plays* are those that have no discoveries or reserves, but which geological analyses indicate may exist. Established plays are categorized further into *ma-ture* and *immature* plays depending on the adequacy of play data for statistical analysis. Mature plays are those plays that have sufficient numbers of discoveries within the discovery sequence so that the *discovery process model* of the PETRIMES assessment procedure is of practical use (Lee and Tzeng, 1989; Lee and Wang, 1990; Lee, 1993). Immature plays do not have a sufficient number of discoveries with established reserves to properly apply the model. Subjective probability play analysis was applied exclusively in this study due to the scarce number of discovered pools with established reserves.

METHOD AND CONTENT

This report incorporates two essential components, geological basin analysis and statistical assessment. Basin analysis fundamentally describes and characterizes the exploration play. Fields and prospects in a play form a natural geological population that can be delimited areally. Once a play is properly defined, a numerical and statistical resource assessment is undertaken using relevant geological data and information for that specific play.

RESOURCE ASSESSMENT PROCEDURE

The analysis of the North Coast area began with the compilation and synthesis of information on regional geology and hydrocarbon occurrence. This included a survey of pertinent publications.

The aim of this data compilation was to initiate basin analysis in order to provide background for the definition of hydrocarbon occurrence models. Play models in the study area were developed by examining the hydrocarbon systems and, when possible, using analogues to extrapolate certain parameters.

Play definition and estimation of reservoir parameters formed the input for a systematic statistical analysis which allowed a quantitative analysis of the undiscovered resource.

GEOLOGICAL PLAY DEFINITION

Definitions of play type and area are essential components of geological basin analysis preceding any numerical resource evaluation procedure. A properly defined play will possess a single population of pools and/or prospects that satisfies the assumption that geological parameters within a play can be approximated by a family of lognormal distributions. Mixed populations derived from improperly defined plays add uncertainty to the resource estimate. Pools and/or prospects in a specific play form a natural geological population characterized by one or more of the following: age, depositional model, structural style, trapping mechanism, geometry, and diagenesis. Prospects or areas within a basin or region can be assigned to specific plays on the basis of a commonality of some or all of these geological elements.

COMPILATION OF PLAY DATA

For conceptual and immature plays, probability distributions of reservoir parameters such as prospect area, reservoir thickness, porosity, trap fill and hydrocarbon fraction are needed. Prospect size can then be calculated using the standard "pool"- size equation. Seismic, well, and outcrop data prove particularly useful in identifying the limits for sizes of prospect area and reservoir thickness as well as porosity limits. Geochemical data are useful in identifying prospective areas as well as the composition of the hydrocarbon accumulations, ie. oil-vs.-gas proneness. Research in similar hydrocarbon-bearing basins is also important in order to provide reasonable constraints on reservoir parameters as well as contribute further information on other aspects of petroleum geology that may prove useful in the study.

SUBJECTIVE PROBABILITY ANALYSIS

There are several methods for estimating the quantity of hydrocarbons that may exist in a play, region or basin (White and Gehman, 1979; Masters, 1984; Rice, 1986; Lee, 1993). Petroleum assessments undertaken by the Geological Survey of Canada are currently based on probabilistic methods (Lee and Wang, 1990) that are developed in the Petroleum Exploration and Resource Evaluation System, PETRIMES (Lee and Tzeng, 1989). The immature and conceptual hydrocarbon plays defined in the North Coast region were analysed by applying a subjective probability approach to the various reservoir parameters. The lognormal option in PETRIMES was used since experience

indicates that geological populations of pool parameters can adequately be represented by lognormal distributions.

Resource assessments in frontier regions use field-size estimates rather than pool-size predictions as derived from mature and immature play analysis in mature basins. A field consists of one or more oil/gas pools or prospects in a single structure or trap. Probability distributions of oil and gas field sizes are computed by combining probability distributions of reservoir parameters, including prospect area, reservoir thickness, porosity, trap fill, hydrocarbon fraction, oil shrinkage and gas expansion.

Probability distributions of oil and gas field sizes are then combined with estimates of numbers of prospects (from seismic and play area mapping) and exploration risks to calculate play potential and to estimate sizes of undiscovered fields.

Exploration risks at a play or prospect level are determined on the basis of the presence or adequacy of geological factors necessary for the formation of petroleum accumulations. Essential factors are reservoir, seal, source rock, timing of hydrocarbon generation, trap closure and preservation. Appropriate marginal probabilities are assigned to each geological parameter. The Yukon coastal plain plays are expected to exist (the low play-level risk of 1.0 was assigned to each play). Within each play, certain prospect-level risks are high and these are assigned appropriate risk factors. Exploration risk is an estimate, incorporating all risk factors, of the percentage of prospects within a play that are expected to contain hydrocarbon accumulations.

Uncertainties in oil and gas play potential and pool-size estimates of a given range of probabilities are necessarily greater than the ranges derived by discovery process analysis used for assessing mature plays. This is due to the nature of frontier assessment results and because few discovered pool sizes are available to constrain sizes of undiscovered accumulations.

REGIONAL GEOLOGY

Two distinct structural and genetic regional geological regimes are represented in northern Yukon. The vast majority of northern Yukon occupies a portion of the Cordilleran Orogen. An area of northeastern Yukon, specifically Peel Plateau, lies on the ancestral North American craton where little Phanerozoic deformation has taken place. This area of ancestral North America is known as the Interior Platform.

There are two major geological provinces lying within the Cordilleran Orogen of northern Yukon. They are separated by the northwest-trending Tintina fault. The area southwest of the fault consists of amalgamated and accreted geological terranes that contain younger, fairly complex assemblages of varying rock-types. The northeastern region, which is part of the morphogeological Frontal Belt, includes a thick assemblage of older sedimentary rocks deposited on the relatively stable ancient North America margin (Hart, 1999).

The western edge of the ancient North America craton extended far out into the ancient Pacific Ocean. This submerged continental shelf of crystalline basement rock is at least 1.7 billion years old and is present throughout northern Yukon beneath both the Interior Platform and the Cordilleran Orogen. These rocks, in part, provided the stable continental platform, upon which sediments, dominantly limestone and sandstone, were deposited over a period of a billion years (Hart, 1999). Shale, sandstone and chert accumulated in basinal regions of deeper water. Thus, the two depositional environments (platform and basin) gave rise to distinct sedimentary packages, dominated by limestone and shale, respectively. These shale and limestone packages are now in fault contact with each other. The Interior Platform amassed thicknesses of between 5 and 25 kilometres of dominant limestone and sandstone. Limestone accumulated during quiescent times in warm, shallow and clear water. The sandstone consists of detritus eroded from the Canadian Shield.



Figure 2. *The major structural elements in the region (from Dixon, et al., 1994).*

A complex folded and faulted terrain, part of the Frontal morphogeological belt of the Cordilleran Foldbelt, underlies the Arctic coastal region of northern Yukon. Structures such as compressional folds and strike-slip faults, formed during the Laramide Orogeny, are underlain by older tectonic features such as the Cache Creek Uplift and Rapid Depression which originated as Mesozoic fault-bounded structures. Other significant older tectonic and structural elements in the region are the Barn and Romanzof uplifts, Blow River and Herschel highs and the Demarcation Subbasin (Fig. 2; Dixon, *et al.*, 1994).

The Rapid Depression (Lane, 1998) is defined on the basis of its Albian fill, with older sediments bounding it to the east, west and south (Dixon, 1996a). To the north, the Blow River High contains Tertiary strata. The Depression is a structurally controlled trough which terminates up-plunge at the upper reaches of the Blow River (Norris, 1997a). Within the trough, more than 5,000 m of Lower Cretaceous turbiditic sandstone and conglomerate were deposited. On the flanks of the trough, this flyschoid succession thins dramatically where it lies unconformably on Ellesmerian units. On top of the flysch, more than 1,200 m of Upper Cretaceous and Tertiary clastics accumulated (Norris, 1997a).

The Cache Creek Uplift underlying southwest Mackenzie Delta is a fault-bounded entity containing rocks ranging in age from at least Middle Devonian to early Tertiary. Middle Devonian or older massive carbonate units in the core of the complex are overlain unconformably by Carboniferous and Permian limestone and sandstone (Norris, 1997a). Deformation in the core appears to be related to the Ellesmerian Orogeny. The Upper Paleozoic rocks are disconformably overlain by Jurassic and Cretaceous shales and sandstones on the northwestern flank of the structure, which, in turn, are covered by lower Tertiary strata (Norris, 1997a).

West of the Rapid Depression lies the Barn Uplift (Fig. 2). It is cored by highly compressed Lower Paleozoic strata, coeval with Road River Formation occurring within the Richardson Mountains to the south. The succession consists of shales, cherts, quartzites, siltstones and limestones (Dyke, 1974). These rocks are overlain unconformably by coarse clastics of the Early Carboniferous Kekiktuk Formation. The rocks in the Uplift are deformed into a series of highly folded, generally steeply dipping panels (Norris, 1997a).

The Romanzof Uplift underlies the British Mountains straddling the Canada-USA border. Isoclinally folded and thrust faulted strata of the Neruokpuk and Road River formations occur within the complex. In Canada, the northern limit of the uplift is in fault contact with the Rapid Depression (Norris, 1997a).

The tectonic elements in the offshore area are principally Late Cretaceous and Tertiary in origin (Dixon, 1996a). The Blow River and Herschel highs are cored by tightly folded Tertiary strata and both have structural elevations higher than the surrounding areas. The Herschel High also contains high-angle reverse faults. A prominent areally large synclinal structure on the south flank of the Herschel High is called the Demarcation Subbasin which is filled with Eocene and younger strata (Dixon, 1996a).

STRATIGRAPHY AND DEPOSITIONAL SETTING

Four major tectono-stratigraphic assemblages are recognized in the Mackenzie Delta–Beaufort Sea region. They include Inuvikian embracing Proterozoic strata, Franklinian encompassing Cambrian to Devonian strata, Ellesmerian which consists of Mississippian to upper Hauterivian rocks, and Brookian comprising upper Hauterivian to modern sediments. Major regional unconformities separate the four assemblages. Significant changes in tectonic regime and activity are designated by these regional unconformities. Another major unconformity is recognized lying beneath Upper Cretaceous strata dividing the Brookian assemblage into an upper and lower subdivision. Most of the oil and gas discovered to date in the Beaufort–Mackenzie area are found in the upper Brookian assemblage (Dixon, *et al.*, 1994).

Figure 3 depicts the stratigraphy and correlative relationships in the Beaufort Sea, Mackenzie Delta and Northern Yukon. Potential reservoir and source rocks in the assessment area are indicated. A schematic cross-section depicting the complex stratigraphic relationships within Upper Jurassic to Paleocene strata beneath the Yukon Arctic coastal plain is depicted in Fig. 4.

INUVIKIAN (PROTEROZOIC)

Interbedded quartzites, argillites, cherts, and limestones of the Hadrynian Neruokpuk Formation are exposed in the core and on the southwest flank of the Romanzof Uplift. The formation is interpreted to also occur beneath the Rapid Depression to the north and east (Norris and Dyke, 1997). Significant discoveries of Paleozoic fossils on the northern flank of the Romanzof Uplift has resulted in reinterpretation of the stratigraphy of the area where previously mapped Precambrian Neruokpuk Formation (Norris, 1981a, b) is now thought to represent strata equivalent to the Road River Group (Lane and Dietrich, 1996). Previous definitions of Neruokpuk strata have included all pre-Mississippian strata in northern Yukon and Alaska (such as, Brosge, *et al.*, 1962; Reed, 1968; Dutro, *et al.*, 1972; Sable, 1977). Lane (1991) argued that Neruokpuk strata should be limited to Proterozoic strata and restricted to the quartzite-dominant unit of that age. These Proterozoic sediments have undergone metamorphism and represent an effective economic basement in the region.

FRANKLINIAN (LOWER PALEOZOIC)

Paleogeographic reconstructions in northern Yukon indicate that Lower Paleozoic strata are generally represented by a basinal facies (Norford, 1997; Lane, 1998). Cambrian to Silurian phyllites, cherts, quartzites and volcanics unconformably overlie the metasedimentary Hadrynian Neruokpuk Formation. These rocks are coeval with Road River Group strata to the south but have sufficient differences in lithologies to warrant separate nomenclature. However, present knowledge of these strata is inadequate for formal nomenclature, so this assemblage of basinal rocks is unnamed (Norford, 1997). There are red and green cherty argillites, shales and phyllites, conglomerates, sandstones and minor limestones in the succession and they are interpreted to have substantial similarities to rocks in Selwyn Basin more than 1,000 km to the south (Cecile, 1988). Thicknesses in northern Yukon are unknown and the rocks are structurally complex. A major unconformity on top of the assemblage locally places Carboniferous, Jurassic or Lower Cretaceous rocks on the Franklinian succession. Lower Paleozoic basinal strata are probably overmature but may retain some potential as a source rock for gas.



Figure 3. *Stratigraphic chart, Beaufort-Mackenzie area (from Dietrich and Dixon, 1997). Potential reservoirs and source rocks in northern Yukon are shown.*

ELLESMERIAN (CARBONIFEROUS TO MIDDLE HAUTERIVIAN)

The lowermost Carboniferous sediments are interbedded chert conglomerates and conglomeratic sandstones of the Kekiktuk Formation that unconformably overlie older strata in the area (Fig. 3; Richards, *et al.*, 1997). Chert, quartz and quartzite clasts are encased in a tight siliceous matrix. This poorly exposed laterally discontinuous unit is exposed in British and Barn mountains and likely occurs through most of northern Yukon. Permian and Carboniferous strata are truncated northeastwards by sub-Triassic and sub-Jurassic unconformities. The Kekiktuk Formation varies in thickness from one to 50 m, averaging about 25 m (Richards, *et al.*, 1997). Lithologies and regional relationships in the Kekiktuk Formation suggest deposition in braided stream and fan delta settings. An underlying regional subaerial erosion surface together with the overlying coastal plain facies of the Kayak Formation infers a continental environment during Kekiktuk deposition. The absence of marine fossils in most of the formation is also indicative of a terrestrial depositional setting. Locally derived, texturally immature sediments with conglomerate channel fills infer the braided stream/fan delta depositional setting (Richards, *et al.*, 1997). Kekiktuk conglomerates are identified as potential reservoir units in the Old Crow plain and on the Alaskan North Slope. Quality of reservoir is unknown beneath the coastal region of the Yukon.

Upper Visean siliciclastics and minor carbonates of the Kayak Formation conformably overlie Kekiktuk rocks. The Kayak Formation consists of basal sandstone with coal seams, a middle shale-dominated unit and an upper interval consisting of shale with subordinate limestone. Thicknesses range from 220 to 375 m in the study area. These clastic formations depict a transgressive, deepening-upward succession where coal-bearing siliciclastics of the basal unit, deposited in shoreline and coastal environments, succeeded fluvial and deltaic sediments in the underlying Kekiktuk Formation (Richards, *et al.*, 1997). Successive shale and carbonate units were deposited in shallow marine to intertidal environments. Coeval carbonate deposition of lowermost Lisburne Group took place in the British Mountains.

In northern Yukon, a widespread Carboniferous carbonate succession consisting of the Alapah and Wahoo formations of the Lisburne Group conformably overlies the



Figure 4. *Schematic cross-section (see Fig. 2 for location) illustrating three clastic assemblages in Upper Jurassic to Paleocene strata of Blow and Kugmallit troughs (after Yorath, 1991).*

Kayak Formation. The Alapah Formation is the thickest and most extensively preserved carbonate unit in northern Yukon. It is more than 1,300 m thick in the western British Mountains and thins to the northeast (Richards, *et al.*, 1997). Lower Alapah carbonates in the British Mountains consist of lime mudstones and wackestones, while skeletal lime grainstones and packstones dominate the upper part. The older carbonates were deposited in restricted to protected shelf environments, while upper Alapah rocks were laid down in a shelf margin setting. Another carbonate unit, called the Wahoo Formation, immediately overlies Alapah carbonates. This formation ranges in thickness from 130 to 225 m in the British Mountains (Richards, *et al.*, 1997). Lime grainstones and packstones dominate this cyclic succession. Deposition probably occurred on protected shelf to shelf margin marine environments. Carboniferous carbonates have been identified as prominent potential reservoir units on the Alaskan North Slope and in Eagle Plain. Dolomitic facies within Lisburne Group carbonates contain significant reserves of hydrocarbons at Prudhoe Bay.

An unconformity separates Carboniferous strata from overlying Permian strata. In the western British Mountains, there is a thin (~200 m), poorly known succession called the Sadlerochit Group that was deposited in a Permian basin which developed north of the Ancestral Aklavik Arch (Fig. 3; Richards, *et al.*, 1997). Local erosional remnants have been preserved in the British Mountains. Shales, sandstone and minor carbonate of the Sadlerochit Group also occur eastward in the Cache Creek Uplift and under southwestern Mackenzie Delta (Dixon, *et al.*, 1994). This unit likely contains a shallow marine facies, including shelf sandstones which may include potential reservoir horizons.

The unconformity-bounded Middle and Upper Triassic Shublik Formation is locally preserved in northwestern Yukon in British and Barn mountains. A nearshore, inner shelf depositional environment is interpreted for Shublik rocks in northern Yukon (Norris, 1997b). This nearshore facies is very heterogeneous with rapid lateral changes in lithology. Basal conglomerate and conglomeratic limestone grade into sandstone, limestone, siltstone, and chert from one exposure to another (Norris, 1997b). Only about 150 m of strata is preserved on the southwest flank of British Mountains. There may be potential for reservoir development among the coarse clastic units in this thin succession in northern Yukon. The Shublik Formation oversteps progressively older formations to the northeast and is erosionally truncated at the sub-Jurassic unconformity. These stratigraphic relationships infer that the southeastward extensions of Permian and Triassic reservoirs at Prudhoe Bay in Alaska do not reach into the Yukon Territory (Norris, 1997b).

Fundamental changes in depositional style and paleogeography took place at the commencement of Jurassic sedimentation. A series of northwestward prograding wedges originated from an arenaceous belt that developed on the southeastern margin of a Jurassic-Neocomian basin ('Brooks-Mackenzie Basin' in Poulton, 1997). These sandstone units grade laterally into shale-dominant units to the northwest and west, away from the exposed North American craton from which they were derived. Interdigitation of arenaceous and argillaceous units in the basin-marginal areas record numerous transgressive and regressive episodes in the overall subsiding shelf regime (Poulton, 1997). In ascending order, the formations within the arenaceous belt in northeastern Yukon and southern Mackenzie Delta are the Bug Creek Group consisting mostly of sandstone, Husky Formation which is shale-dominant, and the Parsons Group which contains the sandstone-dominant Martin Creek Formation, the shales of the McGuire Formation and the sandstone-rich Kamik Formation. In northwestern Yukon, the Bug Creek to McGuire interval is represented singly by the shale succession called the Kingak Formation. The arenaceous belt is interpreted to occur at the basin margin or within

the inner shelf facies, while the argillaceous Kingak succession occupies the outer shelf to the northwest.

Jurassic Husky and Kingak shales have been recognized as potential source rocks in Beaufort-Mackenzie Basin and the Alaskan North Slope. Davies and Poulton (1989) report elevated Thermal Alteration Indices (T.A.I.) and vitrinite reflectance values in outcrop in the western Richardson Mountains. These source rocks have passed through the oil window making them overmature and resulting possibly in the generation of dry gas, which may still be taking place.

BROOKIAN (UPPER HAUTERIVIAN TO EOCENE)

A major regional unconformity occurs at the base of the Upper Hauterivian to Barremian Mount Goodenough Formation. Another major unconformity separates Upper Cretaceous strata from Lower Cretaceous rocks. The mid-Cretaceous unconformity distinguishes two tectono-stratigraphic sequences: a lower assemblage consisting of easterly to southeasterly craton-derived sediments deposited in shoreline and shelf environments, and an upper assemblage derived from the Cordilleran Orogen to the south and southwest. The younger assemblage consists of Upper Cretaceous sediments deposited in a cratonic foreland basin overlain by Tertiary rocks that accumulated in a subsiding continental margin (Dixon, 1997).

Upper Hauterivian and Barremian strata are embodied by the shale-dominant Mount Goodenough Formation. This unit is present throughout northern Yukon and in the subsurface of Mackenzie Delta and the Beaufort Sea. A basal sandstone is locally developed which may contain reservoir facies. This sandstone is especially well-developed adjacent to and upon positive tectonic elements such as the Cache Creek and Romanzof uplifts. Gradationally overlying Mount Goodenough shales in the northern Richardson Mountains are interbedded sandstones and shales of the Rat River Formation. This formation is Barremian to Aptian in age. The basin margin, or shoreline facies, of the Rat River Formation grades laterally to the northwest into the shale-dominant shelf area represented by the Mount Goodenough Formation.

Late Aptian and Albian rocks are designated Albian Flysch in northern Yukon. Thick accumulations of Albian sediment were deposited in grabens and half-grabens such as Blow Trough in the Rapid Depression (Dixon, 1997). Within the Albian Flysch succession, shale is dominant but there are also significant stratal intervals containing sandstone and conglomerate derived from the ancestral Brooks Geanticline to the west. These flysch deposits also underlie the Yukon North Coast to the northwest. The flysch was deposited as sediment gravity-flow deposits in slope, submarine and basinal environments (Dixon, *et al.*, 1994).

The major unconformity separating Lower and Upper Cretaceous strata marks a profound change from the deep-water basins, mud-rich shelf areas and poorly defined foreland basins prominent during Albian time, to the post-Albian emergence of well-defined foreland basins filled with coarse clastics from the Cordilleran Orogen (Dixon, 1997). This unconformity is especially prominent west of Richardson Mountains where shales containing thin beds of bentonite of the Boundary Creek sequence abruptly overlie the Albian Flysch. These shales were deposited within outer shelf to slope environments (Dixon, *et al.*, 1994). The Boundary Creek shales are rich in organic matter, averaging 3 to 5% TOC. The slightly younger Smoking Hills strata present in Mackenzie Delta and the Beaufort Sea are especially rich in organic matter, but are eroded west of Richardson Mountains. The Maastrichtian to Paleocene Fish River Sequence unconformably overlies

the Boundary Creek Sequence in northern Yukon. This unconformity marks a significant shift from cratonic sedimentation to continental margin deposition.

The late Maastrichtian-Tertiary continental margin deposits consist of a series of prograding deltaic complexes. The earliest delta complex was centred over the western Beaufort shelf, whereas subsequent delta centres tended to shift eastward (Dixon, 1997). The earliest delta is defined by the Fish River Sequence in northern Yukon. The younger late Paleocene-early Eocene Aklak sequence, conformably lying on top of Fish River strata, is also centred in the western part of the Beaufort Sea. Both the Fish River and Aklak sequences form an extensive belt of sandstone-rich strata.

Every Tertiary sequence contains a thick succession of interbedded sandstone and shale at the basin margins, distributed in delta plain and delta front deposits. Basinward, the amount of shale increases. The early to middle Eocene Taglu sequence is centred under Richards Island with deltaic deposits extending westward into the shallow waters of Beaufort Shelf in northern Yukon (Dietrich and Dixon, 1997). A thin cover of late Pleistocene-Holocene Shallow Bay basinal shales is present on the northern Yukon continental shelf overlying a Late Miocene unconformity (Dietrich and Dixon, 1997).

STRUCTURAL GEOLOGY

To the west of the Richardson Mountains and extending into western Beaufort Sea, highly deformed strata of the Cordilleran Foldbelt occur (Norris, 1981a, b; Lane and Dietrich, 1996). These compressional and strike-slip structures were formed in late Cretaceous to Tertiary time during the Laramide Orogeny. The arcuate trend of the offshore fold and thrust belt is marked onshore by the eastward to southeastward strike of structures in northwestern Yukon, while further east in Blow Trough and in the Richardson Mountains, the structures trend north-south. Brookian deformation taking place during Eocene time produced these prominent structures which overprint Paleozoic compressional and Jurassic-Cretaceous extensional structures in the study area (Lane and Dietrich, 1996). Tertiary extensional faults are also common throughout the Beaufort-Mackenzie Basin. Older structural features, such as the Rapid Depression and Cache Creek Uplift, probably originated as Mesozoic fault-bounded structures. In the Blow Trough, the rift setting is not apparent due to the lack of preservation of associated structures. Chevron folds and local thrust faults, which may be contiguous with Brookian structures interpreted offshore (Lane, 1988), are present. Blow River and Herschel highs are large-scale tectonic features cored by closely spaced anticlines (Fig. 2).

Rocks on the Yukon North Coast were extensively deformed during Paleocene to early Eocene time (Lane, 1998). Folds and thrust faults observed in Paleocene outcrop (Norris, 1981a), as well as continuity of structures with known Eocene structures offshore, constrain this major deformational event.

PETROLEUM GEOLOGY

EXPLORATION HISTORY

Seismic survey activity in the Mackenzie Delta area in the early 1960s delineated large structures in favourable stratigraphic sections. These early surveys led to the drilling of two dry wells in 1962. Further exploration resulted in the initial discovery of oil in Cretaceous sandstones at Atkinson Point in 1969. In 1970, a major gas find was made in Kamik sands at Parsons Lake and in the same year oil was discovered in carbonates at Mayogiak. In 1977, the focus of exploration switched offshore to Tertiary targets. In the Beaufort-Mackenzie region, 53 oil and gas discoveries, both onshore and offshore, have been made. Forty-four of these discoveries occur in the Tertiary basin. A total of 247 wells has been completed to date in the Beaufort-Mackenzie region. Drilling on the Yukon North Coast to the west has been very limited. Three wells were completed showing no hydrocarbons and limited reservoir potential.

RESERVOIRS

CARBONIFEROUS

A gas discovery was made in Carboniferous carbonates at Unak L-28 in west-central Mackenzie Delta. This discovery exhibits an instance when the generally poor primary porosity development of Carboniferous carbonates was enhanced by fracture development in the south delta. Dolomitized facies and leached zones adjacent to unconformities occasionally produce porous horizons in marine Upper Carboniferous Alapah and Wahoo carbonate formations of the Lisburne Group. There have been discoveries of hydrocarbons in equivalent rocks on the Alaskan North Slope and in southern Eagle Plain (Hamblin, 1990). Porosity development in the subsurface of the Yukon is unknown but surface exposures indicate that primary porosity is generally poor. Secondary porosity development may be significant but patchy.

A possible reservoir interval that may be present in the Yukon Arctic coastal region is the Lower Carboniferous Kekiktuk Formation (Fig. 3). This formation, containing quartz, chert, quartzite and granitic clasts, generally occurs in a tight siliceous matrix. Even though porosity development in this unit is commonly poor in the subsurface, leaching has produced significant reservoirs in Alaska, specifically at the Endicott Field (Craig, *et al.*, 1985). Primary porosity is very low or absent, but secondary, patchy porosity development may occur.

PERMO-TRIASSIC

Permian and Middle to Upper Triassic potential reservoir strata, represented by Echooka sandstones of the Sadlerochit Group and coarse clastics of the Shublik Formation, respectively, exhibit good porosity development in Eagle Plain and the Alaskan North Slope. Paleogeographic reconstructions (Bamber and Waterhouse, 1971) indicate that these potential reservoir units may occur beneath the North Coast of the Yukon. Young (1981) considers the Sadlerochit Formation to be a prime prospect for future hydrocarbon discoveries in northern Yukon. Primary and secondary porosity may develop in both conglomerate and sandstone units.

JURASSIC

Northwest-prograding clastic wedges of the Jurassic arenaceous belt, located in northern Richardson Mountains and in the subsurface beneath southern Mackenzie Delta, have potential as reservoir rock. Bug Creek sandstones in the area retain fair potential as reservoir-quality rock, although these rocks are typically tight and no discoveries have been made. Porosity exceeds 10% in some of the cleaner facies but there is low permeability in the coarse clastics (Northern Oil and Gas Directorate, 1995a). To the northwest along the Yukon North Coast, the Jurassic Brooks-Mackenzie Basin deepens, whereupon finer-grained clastics are deposited, substantially reducing reservoir potential.

CRETACEOUS

Hauterivian Kamik sandstones of the Parsons Group are excellent potential reservoir horizons in the Mackenzie Delta. Both oil and gas have been recovered from this thick, extensively developed unit. Average porosity at Parsons Lake is 15%. Kamik sands in the Yukon Territory occupy a sandy inner to mid-shelf belt which likely has inferior reservoir qualities compared to the interpreted beach and inner shelf environment of the Delta to the east (Dixon, 1986).

There is a thin basal sandstone with local lenses of coarser material in the Mount Goodenough Formation that seems to be fairly persistent throughout the area. A minor oil discovery was made in this interval proving the reservoir potential in this strata. Late Barremian to Aptian Rat River sands, which are interpreted as representing a regressive pulse, contain excellent potential reservoir horizons in western Mackenzie Delta, as indicated by the gas discovery at Unak L-28 (Dixon, 1986).

There is little or no reservoir potential in Upper Cretaceous strata. Albian sedimentation is dominated by fine-grained deep-water clastics deposited during increased rifting and subsidence. Blow Trough, a depositional precursor of the Rapid Depression, extended over a large area of northern Yukon and provided a major site for Albian deposition. Cenomanian to Turonian Boundary Creek rocks also comprise a shale-rich basinal sequence, indicating poor reservoir potential.

TERTIARY

Most Tertiary reservoirs occur offshore but there is some potential for reservoir occurrence along the Arctic coastal plain, if trapping conditions are favourable. Oil staining occurs in similar Tertiary sands in Alaska. Sandstone members in the Maastrichtian-Lower Paleocene Fish River Sequence are potential reservoirs. The Fish River deltaic complex seems to be centred over the western part of the Mackenzie Delta, specifically in the Mackenzie Bay area (Dixon and Dietrich, 1996). The bulk of the sandstone-dominant part of the sequence underlies the western Beaufort shelf which remains to be tested by drillholes. Deltaic rocks of the overlying Paleocene-Lower Eocene Aklak Sequence are recognized in the western Beaufort Sea shelf area. Oil and gas have been recovered from this sequence further east in the Mackenzie Delta area. Delta-front deposits of the Aklak sequence at Adlartok in Yukon offshore waters to the north are characterized by highly variable porosity and permeability due to grain size. Coarser beds exhibit a porosity range of 20 to 24% while finer-grained laminated units vary from 15 to 18% (Northern Oil and Gas Directorate, 1995b).

SEALS

The laterally and vertically extensive Jurassic and Lower Cretaceous Kingak shale provides excellent top seal for Upper Paleozoic and Triassic reservoir strata. Additional Lower Cretaceous marine shale-rich units such as the Mount Goodenough Formation and the Albian Flysch are effective top seals for potential Cretaceous sandstone reservoirs. Another efficient top seal for all pre-Tertiary reservoirs in northern Yukon is the widespread Boundary Creek shale-rich sequence. Intraformational shales form localized top and lateral seals for Tertiary reservoirs.

Numerous unconformities in the study area seal underlying reservoir units by juxtaposing permeable and impermeable formations. Young (1981) indicated that the pre- Upper Triassic unconformity in northern Yukon may effectively seal older strata in broad areas and numerous updip traps may be developed along the edges of pre-Upper Triassic structures.

TRAPS

A variety of structural, stratigraphic and combination traps occur within Phanerozoic sedimentary strata throughout the region. Pre-Laramide traps are more favourable for accumulating hydrocarbons since the primary episode of hydrocarbon generation probably took place in Mesozoic and Tertiary time when maximum load was imposed. In southern Mackenzie Delta, trap types are dominantly structural where closure often occurs against normal faults. Further west, thrust faults complicate the pre-existing pattern of tilted fault blocks. Along the Yukon North Coast, numerous structural features have been recognized. There are broad anticlinal features, domes, and normal and reverse faults, along with horst and graben structures. In Herschel and Blow River Highs, folds are closely spaced and have large amplitudes. Reverse fault traps are also present. Unconformity-related combination traps occur throughout the assessment area.

SOURCE ROCKS

Numerous potential source rocks for oil and gas have been identified in the Beaufort-Mackenzie region. Upper Cretaceous mudstones and shales of the Boundary Creek Sequence are considered to be significant source rock for oil. The sequence is rich in organic matter with TOC ranging from 3 to 5%. The richer oil-prone Smoking Hills Sequence (2 to 10% TOC) has been eroded in western Mackenzie Delta and does not appear to extend into the Yukon Territory (Dixon, 1996b). Smoking Hills-sourced oil constitutes the Kugpik O-13 Cretaceous discovery and there are reasonable expectations that Smoking Hills-sourced oils may have migrated further west into Yukon Territory.

The Jurassic Kingak and Husky shale succession is a significant gas-prone source rock in the area. Langhus (1980) indicates that the terrestrially dominant organic matter present in these rocks is consistent with a gas-prone source rock which may have provided the charge for gas accumulations at Parsons Lake. At Unak L-28, gas was recovered from both Carboniferous carbonates and the Lower Cretaceous Rat River Formation. It is believed that gas for both of these accumulations is derived from the Jurassic-Cretaceous shale succession.

Oil and gas found in Tertiary deltaic reservoirs in the central Beaufort region are most likely derived from the basal portion of the Richards shale sequence (Brooks, 1986). However, to the west, specifically at Adlartok, oil geochemistry indicates that

the Richards-type biomarker is not present. Oil at Adlartok may have been derived from Paleocene shales. Organic matter is not particularly concentrated in these Tertiary prodelta depositional environments (infrequently exceeding 2% TOC), but the source rock intervals are thick. Tertiary-derived oils are generally good quality (ranging from 25 to 35° API). There are also significant condensate volumes associated with some of the onshore gas discoveries.

Argillites and shales of the unnamed Cambrian to Silurian succession coeval with Road River Formation basinal shales may be locally organic-rich in the Arctic coastal plain area. Link, *et al.*, (1989), however, rated the overall source rock potential of this formation in northern Yukon as poor. Hydrocarbons generated by Road River-equivalent strata were available for migration during Devonian to Carboniferous time well before the significant Laramide deformational event. Lower Paleozoic strata in this area are overmature and possibly supermature, but may retain limited potential for gas generation. In southern Mackenzie Delta, the Lower Paleozoic shales have moderate organic quality but poor source potential due to low TOC (Link, *et al.*, 1989).

Carboniferous shales are reported to be an important source rock in surrounding areas, specifically beneath the Alaskan North Slope and in Eagle Plain (Craig, *et al.*, 1985; Link, *et al.*, 1989; Utting, 1989). In the Alaskan North Slope area, Kayak Formation shales are lean and gas-prone with some potential for oil generation. In Eagle Plain, the equivalent Carboniferous shales are characterized as fair to good source rocks for gas with some oil potential (Link, *et al.*, 1989). Comparable shales found in the Yukon North Coast area, therefore, are expected to have at least fair potential to generate gas if maturities are equivalent. Thermal maturities of these rocks are likely high and probably well beyond the oil window. No Kayak samples have reported vitrinite reflectance values of less than 2.5% and Thermal Alteration Indices (TAIs) range from 3+ to 5 in British and Barn mountains south of the North Coast (Cameron, *et al.*, 1986; Utting, 1989).

Although the Late Triassic Shublik Formation is considered to be a significant oil source at Prudhoe Bay (Claypool and Magoon, 1985), the high-energy carbonate facies, occurring only in extreme northwestern Yukon, is unlikely to be a source rock of comparable quality.

TIMING OF HYDROCARBON GENERATION

There are numerous Tertiary organic-rich horizons that exhibit hydrocarbon source rock potential in the Beaufort-Mackenzie area such as occurrences in Boundary Creek, Smoking Hills and Richards sequences. Maximum burial probably occurred during middle Eocene time when hydrocarbons were likely generated. Several stages of Tertiary contractional deformation are identified in Tertiary sediments in nearshore areas of the Yukon North Coast. Maturity of organic-rich Tertiary sediments was likely reached contemporaneously with later stages of the Laramide orogeny and previous to subsequent pulses of tectonic deformation. At least some structures formed during the early stages of the Laramide deformation were available for accumulating hydrocarbons.

Modelling by Link and Bustin (1989) of Paleozoic and Mesozoic source rocks in northern Yukon indicate that they passed through the 'oil window' before the end of Mesozoic time. This implies that the most effective trapping configurations for pre-Tertiary hydrocarbons are structures formed previous to the Late Cretaceous–Tertiary Laramide orogeny, when hydrocarbons generated during the period of active oil migration were trapped in pre-Mesozoic traps. Seeking Paleozoic and Mesozoic targets

based on Laramide anticlinal structures may not represent the most efficient exploration strategy for this region. Pre-Tertiary traps and reservoirs are more favourable sites for accumulation of hydrocarbons generated during late Paleozoic to Mesozoic times.

Stratigraphically trapped Jurassic sandstone and Paleozoic sandstone and carbonate reservoirs on the Yukon North Coast are located updip of the Tertiary hydrocarbon kitchen buried in the deep sedimentary pile to the north beneath the Tertiary deltas.

HYDROCARBON SHOWS

The most direct indication of oil and gas potential in a frontier region are hydrocarbon pools and shows. Hydrocarbon shows and recoveries have been made in Paleozoic carbonates, Lower Cretaceous sandstones and Tertiary sandstones in the Beaufort-Mackenzie region. All Paleozoic and Lower Cretaceous reservoirs are found in southern Mackenzie Delta and along the Arctic coastal plain, and Tertiary discoveries are mostly concentrated in the central Beaufort-Mackenzie Basin. West Beaufort Sea shows potential with the discovery of oil at Adlartok in Paleocene and Eocene sands.

Limited discoveries have been made in Paleozoic carbonate reservoirs: a total of three hydrocarbon accumulations throughout the basin. Oil was discovered in the eastern portion of the basin, while at Unak L-28 in southwestern Mackenzie Delta, gas was recovered from Paleozoic reservoirs. The Unak trap features closure against a thrust fault. The gas is likely derived from overlying Jurassic and Cretaceous shale.

Oil and gas has also been recovered from Lower Cretaceous sandstones throughout the onshore portion of the Beaufort-Mackenzie Basin. Oil is found in the Parsons Group sands, the Atkinson Point Formation and the Mount Goodenough Formation. In western Mackenzie Delta, the Kugpik O-13 well produced oil from the Parsons Group sandstone reservoir. Smoking Hills shales are interpreted to contain oil-prone organic matter for Cretaceous reservoirs. Gas was found in the Lower Cretaceous Rat River sandstone at Unak L-28. Source for Cretaceous Unak gas is unknown although the thick Jurassic-Cretaceous shale succession certainly has potential.

Most of the hydrocarbon discoveries in the basin occur in Tertiary strata. Widespread and thick deltaic coarse clastic deposits provide ample potential reservoir horizons. Interbedded and widespread organic-rich shale layers provide significant oil and gas source material. Oil and gas have been recovered from the Fish River, Aklak, Taglu, Kugmallit and Mackenzie Bay sequences. There seems to be a general trend of oil-prone hydrocarbons occurring basinward, although numerous deviations off this trend have been observed from discoveries made. Richards shales and organic-rich horizons in Paleocene shales are possible oil sources. The Tertiary succession is dominated by Type III terrestrial organic matter. Conventionally, this type of organic matter produces gas. In Beaufort-Mackenzie, large volumes of oil are encountered in Tertiary reservoirs as well as significant quantities of gas. Snowdon and Powell (1979) suggested that these oils were generated from resin-rich terrestrial organic matter. The resinous matter can also generate hydrocarbons at lower levels of organic maturity which is consistent with the interpreted thermal regime (Dixon, *et al.*, 1994). No Tertiary discoveries have been made in the Yukon portion of the North Coast study area. Oil was recovered from Paleocene and Eocene reservoirs at Adlartok in Yukon waters north of the study area.

HYDROCARBON ASSESSMENT

The Yukon North Coast petroleum resource assessment was undertaken in order to generate quantitative estimates of total oil and gas potential and possible sizes of undiscovered fields. Hydrocarbon assessments of basins or regions are usually based on analyses of a number of exploration plays. In the North Coast region, six exploration plays were defined based on various petroleum geological considerations such as structural style, dominant reservoir lithology and thermal maturity. Three immature and three conceptual oil and gas plays were identified in the study area. The conceptual and immature plays had sufficient information to attempt a statistical analysis in order to obtain ranges of estimates of resource potential and sizes of individual undiscovered fields.

Table 1. *Hydrocarbon potential in Yukon North Coast assessment region.*

Play Name	Expected number of fields (mean)	Median play potential (million m^3)	Remaining median play potential (million m^3)	Median of largest field size (million m^3)	Median of largest remaining field size (million m^3)
Oil Plays					
South Delta–Mesozoic	4	2.8	2.4	1.5	0.7
Herschel	5	34.6	34.6	15.5	15.5
All oil plays	9	39.8*			
Gas Plays					
South Delta–Mesozoic	8	2,040	1,217	750	391
South Delta–Paleozoic	15	6,661	6,370	1,720	1,720
Herschel	5	16,095	16,095	7,146	7,146
Yukon Coastal Plain	3	8,463	8,463	5,998	5,998
All gas plays	31	38,565*			
Condensate volumes	**Average liquid content (m^3/million m^3) (million m^3)**	**Median play potential (million m^3)**	**Remaining median play potential**		
South Delta–Paleozoic	10.7	0.71	0.68		

* The total oil and gas potential are statistically derived. They are retrieved from the basin empirical distribution in Appendix 2. The median total of basin potential is the value at 50%.

Table 2. *Hydrocarbon play potential in Yukon Territory*

Play Name	Remaining median play potential (in-place) (million m^3)	Median of largest remaining field size (in-place) (million m^3)	Play potential in Yukon (Scenario 1) (million m^3)	Play potential in Yukon (Scenario 2) (million m^3)
Oil Plays				
South Delta–Mesozoic	2.4	0.7	0.74	0.04
Herschel	34.6	15.5	31	15.5
Gas Plays				
South Delta–Mesozoic	1,217	391	412	21
South Delta–Paleozoic	6,370	1,720	1,836	116
Herschel	16,095	7,146	14,395	7,249
Yukon Coastal Plain	8,463	5,998	8,463	N/A

Scenario 1: Largest undiscovered field is assumed to occur in the Yukon Territory.
Scenario 2: Largest undiscovered field is assumed to occur outside the Yukon Territory.

SOUTH DELTA–MESOZOIC OIL AND GAS PLAY

HYDROCARBON PLAYS

Play definition

The South Delta-Mesozoic play encompasses all oil and gas prospects occupying structural and stratigraphic traps involving reservoirs in Jurassic and Lower Cretaceous sandstones underlying the southern part of the Mackenzie Delta (Figs. 5, 6). The play area is bounded to the east by the Caribou Hills, west by the northern Richardson Mountains, south by the Eskimo Lakes Fault Zone and north by the Kugpik area in the Mackenzie Delta. Most of the play area occurs in the Northwest Territories with the remainder occupying a small portion of northeastern Yukon North Coast (Fig. 5). About 2.5% of the play area occurs in the Yukon Territory.

Geology

A thick succession of interlayered shale- and sandstone-dominant strata of Jurassic to Early Cretaceous age underlies the play area. Thickness of the prospect succession varies from 10 to 1500 m and occurs at depths ranging from 10 to 3,750 m. Numerous potential reservoirs are present in the play, such as Bug Creek Group, Parsons Group, basal sandstone of the Mount Goodenough Formation and the Rat River Formation (Fig. 3).

There are two major unconformities within the succession that prove crucial in delineating reservoir units in the play area due to the truncation of reservoir horizons. At the base of the Mount Goodenough Formation, a major unconformity signifies an erosional episode that has removed substantial strata of the Parsons Group in the south especially adjacent to the Eskimo Lakes Fault Zone and next to the Cache Creek Complex (Fig. 2). The other unconformity separating Upper from Lower Cretaceous strata



Figure 5. *South Delta-Mesozoic oil and gas plays (from Dixon, et al., 1988). Wells discussed in report are shown (1-Kugpik O-13, 2-Unak L-28, 3-Ulu A-35).*

has removed Hauterivian to Albian strata in the northern part of the play area, placing Upper Cretaceous strata directly on top of Parsons Group sands.

Sandstones of the Lower to Middle Jurassic Bug Creek Group have fair reservoir potential in the area. These coarse units were deposited as shelf sandstones prior to rifting and subsidence. Porosity exceeds 10% in the cleaner facies, but permeability is low. The distribution of the Group is profoundly affected by the two unconformities and there is also a depositional pinchout to the east.

The thickest and most widespread potential reservoir interval is the Berriasian to Hauterivian Parsons Group. Formations within the Group are Martin Creek, McGuire and Kamik units. Originally, this Group underlay the whole play area but erosion has locally removed strata and modified thickness trends. One oil pool and gas show were discovered in Kamik sands in the Kugpik O-13 well and oil was recovered during a drill-stem test in a Kamik reservoir unit in the Unak L-28 well.

The late Barremian to early Aptian Rat River Formation consists of a set of coarsening-upward sequences characteristic of marine bars or shoals. Prospective strata are limited to the northwestern portion of the play area due to erosion and facies change to argillaceous rocks further south and northeast. One gas pool at Unak L-28 has been discovered in the Rat River Formation.

The Albian Flysch is not generally regarded as containing significant reservoir-quality sands but a gas show was noted in these strata in the Fish River B-60 well.

Dominant trap types in the play are structural with closure against faults representing the most common configuration (Fig. 6). Normal fault traps are important but Unak L-28 shows repeated Mesozoic units which imply thrust faulting complicating the arrangement of tilted fault blocks. Thrust faults are only present in the northwestern part of the play area although seismic reflection data of poor quality in other parts of the play area may mask thrust faults and their related closures (Dixon, *et al.*, 1994).

The Upper Cretaceous Boundary Creek and Smoking Hills sequences are the richest oil-prone source rocks in the region. Erosion, however, has removed this rock from southern parts of the play area. Total organic carbon (TOC) in Boundary Creek/Smoking Hills



Figure 6. *Schematic sketch of the trap-types in the South Delta–Mesozoic oil and gas plays (from Dixon, et al., 1994).*



Figure 7. *Rock-eval/TOC analyses for Ulu A-35 well (from Snowdon, 1990).*

sequences range from 2 to 10%. Another potential oil source is found in organic-rich marine shales of the Albian Arctic Red Formation which are coeval with the Albian Flysch succession (Fig. 3). These rocks are immature due to shallow burial in the area. Terrestrially derived organic matter, generally indicative of gas-prone source material, predominate in Jurassic and Lower Cretaceous shales. The Ulu A-35 well represents one of the 14 wells penetrating the succession. A Rock-eval/TOC analysis was completed on this well (Fig. 7; Snowdon, 1990; 1996). The well was collared in the Plio-Pleistocene Iperk Sequence and penetrated the lower Tertiary sequence, specifically Aklak and Fish River sequences, as well as Cretaceous, Jurassic and Permian strata. The Tertiary succession is marginally mature to mature with the top of the conventional oil window occurring at a depth of 1,370 m. Basal Tertiary, Cretaceous, Jurassic and upper Permian strata occur within the conventional oil window. Depths greater than 3,500 m have samples that are overmature. Hydrogen indices (HI) and TOC values decline accordingly with increases in maturity. The Smoking Hills and Boundary Creek TOC values in the well are somewhat lower than values obtained elsewhere in the basin (averages of 2.33



Figure 8. *Estimate of in-place oil potential of the South Delta-Mesozoic play. Median value of probabilistic assessment is 2.8 million m³ of in-place oil distributed in 4 fields.*

and 1.95%, respectively). Husky Formation TOC average values in the well (1.95%) are comparable to Smoking Hills and Boundary Creek sequence total values. Hydrogen index values in Husky shales are only slightly lower than Cretaceous units (Snowdon, 1996).

Exploration risks

All of the Yukon North Coast immature and conceptual plays are inferred to exist (implicated by a play-level marginal probability of 1.0). However, within each play, geological risk factors at the prospect level are evaluated in order to derive the exploration risk for the entire play. Significant prospect-level risks associated with this play are presence of closure, and adequate porosity, source and preservation (Appendix 1). Closure risk is related to significant difficulty in mapping and interpreting closures against thrust faults in the Unak area. Well-cemented sandstones are common in the play making porosity generally low and increasing the risk for reservoir facies in certain prospects. High risk was assigned to source since potential source material may be eroded and



Figure 9. *Estimate of in-place gas potential of the South Delta–Mesozoic play. Median value of probabilistic assessment is 2.0 billion m³ of in-place gas distributed in 8 fields.*

removed over large portions of the play area. Cretaceous erosional events may also remove reservoir facies in parts increasing the risk of adequate preservation in the play.

On the other hand, low risk, or high marginal probability, was assigned to seal and timing (Appendix 1). Thick shale successions separate sandy formations providing effective top seal for most of the prospects. Most structures, particularly closures against normal faults, seem to have formed previous to the main episode of hydrocarbon generation when maximum burial was achieved, thus, providing structural traps before hydrocarbon migration took place. Younger thrust fault structures in the north, however, may have post-dated the main episode of hydrocarbon generation.

Play potential

The South Delta–Mesozoic play has estimated in-place median oil and gas resource values of 2.8 million m^3 and 2.0 billion m^3, respectively (50 percentile values in Figs. 8, 9; Table 1). If the 95 and 5 upper percentiles representing the range of expected resource are specified,



Figure 10. *Field-size-by-rank plot of the South Delta–Mesozoic oil play. Median value of the largest predicted field size is 2.4 million m^3 of in-place oil.*

then there is a 90% chance that the resource potential of oil resides within the range of 0.2 to 10.5 million m^3 in-place (Fig. 8). The range for in-place natural gas of expected resource potential is 0.5 to 5.7 billion m^3 in-place (Fig. 9). In the Kugpik O-13 well, one oil pool has been discovered. It has an in-place volume of 6.7 million m^3. The in-place median play potential conditioned on this discovery is reduced to 2.4 million m^3. Similarly, one gas pool was discovered in the play at Unak L-28, containing an in-place volume of 1,578 million m^3. The conditioned median potential for gas resource is 1.2 billion m^3.

The mean value of the number of predicted fields for the oil play is four. The largest field is expected to contain 1.5 million m^3 of oil (median value of first box in Fig. 10, Table 1). The Kugpik O-13 pool, with a volume of 6.7 million m^3, matches with the fifth upper percentile of the largest predicted field size (see Appendix 2 for computation outputs). Therefore, the largest pool or field has already been discovered and three more fields are predicted. The median of the largest remaining field size is 0.7 million m^3 in-place (median value of second box in Fig. 10; Appendix 2). No fields with volumes greater than 160 million m^3 of oil are predicted to occur in this clastic play. This threshold value,



Figure 11. *Field-size-by-rank plot of the South Delta–Mesozoic gas play. Median value of the largest predicted field size is 750 million m3 of in-place gas.*

160 million m^3 (1 billion bbls), is an arbitrary in-place oil volume defined as the minimum individual field size needed in a specified play on a stand alone basis in a frontier region of Canada that may spark interest among explorationists for initiation of exploration activities in that region.

In the gas play, the expected number of natural gas fields is 8. The median in-place size of the largest predicted field size is 750 million m^3 of gas (Fig. 11, Table 1). The discovered gas pool at Unak L-28 with a volume of 1,578 million m^3 lies well within the 95/5 upper percentile range for the largest predicted field size. The largest gas field, therefore, has been discovered in the play and seven smaller fields are predicted. The largest remaining field has a median estimate of 391 million m^3 (Fig. 11, Appendix 2). A threshold volume of 3,000 million m^3 or 100 BCF has been arbitrarily defined as a minimum field size needed to spark sufficient interest for exploration of a single play in a frontier region. No fields of this minimum size are predicted for this play.

The method employed in this study where oil and gas assessments of the same play are run separately, does not reveal how the oil and gas is distributed among the twelve predicted fields of both plays. It is not known what proportion of individual oil and gas accumulations occur as saturated or undersaturated pools in the same prospect or as separate pools in different prospects.

Hydrocarbon plays in this assessment area often occupy areas spanning political boundaries such as the Yukon/Northwest Territories and Yukon/Alaska borders. If it can be assumed that the hydrocarbon resource is evenly distributed throughout the total play area, the proportion of resource occurring in the Yukon can be estimated by comparing play areas among the various political jurisdictions. The location of the largest remaining field cannot be determined, so two scenarios are proposed; scenario 1 where the largest field is assumed to occur in the Yukon, and scenario 2 where the largest field is present outside the Yukon Territory. Table 2 itemizes the diminished oil and gas potential for the North Coast region in the Yukon Territory under the two proposed scenarios. Only about 2.5% of the play area of the South Delta–Mesozoic oil and gas play is located in the Yukon (Fig. 5), so the amount of oil and gas predicted if the largest undiscovered field is assumed to occur in the Yukon is 0.74 million m^3 and 412 million m^3, respectively (Scenario 1). If, however, the largest remaining field is assumed to reside in the Northwest Territories, then the potential is reduced substantially to 0.04 million and 21 million m^3 of oil and gas, respectively (Scenario 2; Table 2).

SOUTH DELTA–PALEOZOIC GAS PLAY

HYDROCARBON PLAYS

Play definition

This exploration play contains all structures and prospects occurring in Upper Carboniferous to Permian reservoir strata in the southern Mackenzie Delta (Fig. 12). The play area coincides with the South Delta–Mesozoic oil and gas play (Figs. 5, 12). Gas is the principal hydrocarbon. Only 2.5% of the play area occurs in the Yukon Territory.

Geology

Prospective targets in this Upper Paleozoic succession are dolomitic facies and leached zones adjacent to unconformities within marine limestones of the Upper Carboniferous Lisburne Group as well as local coarser strata among interbedded sandstones and shales of the Permian Sadlerochit Group. The prospect succession may attain thicknesses up to 500 m and it occurs at depths ranging from 1,200 to 4,000 m. Ten wells have penetrated Paleozoic strata in the play area resulting in one discovered gas pool at Unak L-28 occurring in Lisburne carbonates. Condensate was also recovered from Carboniferous strata in the same well. Porosity and permeability of the Upper Paleozoic succession is generally poor, but sufficient secondary porosity development resulting from fracturing has created some reservoir potential.

Structural traps are likely dominant, with closure against normal faults most common (Fig. 13). In the northwestern part of the play area including the area in the Yukon Territory, thrust faulting is present which complicates the tilted fault block pattern. Unconformity-related stratigraphic traps are also likely present. Marine shales in Jurassic and Lower Cretaceous strata provide effective top seal for these traps in Paleozoic strata. Most of these structures developed during the Laramide orogeny. Hydrocarbon



Figure 12. *South Delta–Paleozoic gas play (from Dixon, et al., 1988). Wells discussed in report are shown (1–Unak L-28, 2–Ulu A-35).*

generation likely occurred during the latter stages of the orogeny when maximum burial was achieved. Therefore, most structures were formed previous to or contemporaneous with the main episode of hydrocarbon generation. Younger thrust-related structures seem to have developed subsequent to the main phase of hydrocarbon generation. Renewed burial of organic-rich sediments by thrusting episodes may have induced a secondary phase of thermogenic gas generation.

Beneath the Mackenzie Delta, Upper Paleozoic source rocks have not been identified. The juxtaposition of Paleozoic reservoir against Mesozoic organic-rich shales by faulting produces favourable trapping configurations. In the Ulu A-35 well (Fig. 7), Rock-eval analyses indicate that Upper Permian strata lie within the lower portion of the conventional oil window, meaning that organic matter occurring in Lower Permian strata is overmature (Snowdon, 1990, 1996). This implies that any hydrocarbon generated from Paleozoic source rock in the area is likely dry gas only. In the well, most organic matter seems to be terrestrially derived (Type III; Fig. 7). Type III organic matter usually indicates gas-prone strata. HI and TOC values also decrease with depth as maturity increases. Little or no oil is expected in this play.

Exploration risks

Heightened risk factors integrated into the analysis of this exploration play are presence of closure and adequacy of porosity (Appendix 1). Porosity development is generally poor and inadequate reservoir quality may be widespread. Closure integrity may be affected by erosion. Adequate seal, timing, preservation and source are expected for most prospects in this play.

Play potential

This play has an estimated median resource potential of 6.7 billion m^3 of in-place natural gas (Fig. 14; Table 1). The range of estimates for resource potential is 1.7 to 16.5 billion m^3 in-place. The expected number of gas fields in the play is 15 (mean value) with the largest field having a volume of 1.7 billion m^3 (Fig. 15; Table 1). No fields are expected with volumes greater than three billion m^3 in this play (Appendix 2). One



Figure 13. *Schematic sketch of the trap-types in the South Delta-Paleozoic gas play (from Dixon, et al., 1994).*

gas pool was discovered at Unak L-28 in the Northwest Territories having an initial in-place volume of 311 million m^3. This pool was discovered in Carboniferous carbonates. The expected median estimate of the play potential conditioned on this discovery is 6.4 billion m^3 (Appendix 2). The gas discovery matches closely with the median volume retrieved for the seventh largest predicted field size. Six larger field sizes, therefore, are predicted to occur in this play. Condensate was also recovered at Unak in Carboniferous carbonates. An average liquid content of 10.7 cubic metres of condensate per million cubic metres of gas in the Unak gas pool gives a median potential of 0.71 million m^3 of condensate in the play. Remaining median potential, discounting the Unak discovery, is 0.68 million m^3 (Table 1).

Only 2.5% of the play area is defined in the Yukon Territory (Fig. 12). If the largest undiscovered field is assumed to occur within this small area, then the play potential in the Yukon is 1.8 billion m^3 (Scenario 1; Table 2). If the undiscovered field remains to be found in Northwest Territories (which is much more likely), the play potential in the Yukon is much reduced, to 116 million m^3 in-place (Scenario 2; Table 2).



Figure 14. *Estimate of in-place gas potential of the South Delta–Paleozoic play. Median value of probabilistic assessment is 6.7 billion m^3 of in-place gas distributed in 15 fields.*

HERSCHEL OIL AND GAS PLAY

HYDROCARBON PLAYS

Play definition

The Herschel play includes all Paleocene to Early Eocene Fish River and Aklak sequence oil and gas prospects present in Herschel and Blow River highs (Figs. 2, 16). These two structural complexes, containing highly deformed Tertiary strata, occur beneath the shallow waters of the southwestern Beaufort shelf. The Blow River High extends onshore beneath the coastal plain near the Yukon-Northwest Territory boundary. The Herschel High continues westwards into Alaska offshore waters. Much of the play area occurs in the Yukon Territory (about 81%) (Fig. 16).

Geology

Paleocene and Early Eocene strata are intensely folded and faulted within cores of two structurally elevated complexes that form part of the west Beaufort Tertiary fold belt. High-amplitude, closely spaced folds and reverse faults represent the most common structural trap type in the play (Fig. 17). Sandstones may subcrop unconformities forming small stratigraphic traps. Within these deformed Tertiary strata, potential reservoir units include delta-front and delta plain sandstones. Thickness of this succession ranges from 5,000 to 7,000 m in the play area. Depths of prospects vary from 1,500 to 8,000 m. Deep erosion of sub-middle Eocene strata has occurred over the crests of these highs. No wells and no hydrocarbon shows are recorded in the play.

Several stages of Tertiary contractional deformation have been identified within the play area. A major tectonic pulse, affecting lower Tertiary strata, commenced in Early Eocene and culminated in Middle Eocene time. Folds and thrust faults were formed at this time. Post middle Eocene deformation significantly overprinted this earlier stage



Figure 16. *Herschel oil and gas plays (from Dixon, et al., 1988).*

of deformation. Identifiable deformational stages occur in late Eocene, Oligocene and late Miocene time. Maturation of potential hydrocarbon sources was reached prior to completion of structural deformation. Maximum burial likely took place at about 85 Ma, previous to the start of Laramide deformation at 78 Ma. Migration of hydrocarbons was initially controlled by depositional geometries. Hydrocarbons trapped during these initial stages may have subsequently re-migrated into anticlinal and fault-related traps following deformational episodes.

Oil and gas source rocks are found in Upper Cretaceous Boundary Creek and upper Eocene Richards shales. Oil-prone strata are also found in the lower Fish River sequence. The organic matter is dominantly Type III terrestrial-dominated material. The organic matter is mature, however, TOC values are generally quite low with many intervals containing only 1-2% TOC. Conventionally, Type III organic matter generates gas but the oils interpreted to be derived from these organic-rich horizons may be produced from resin-rich terrestrial organic matter.

Exploration risks

Significant risk factors attributed to the Herschel play are the presence of closure and adequate seal and source (Appendix 1). Lack of seal and closures are related to the deep erosion of Fish River and Reindeer strata over the structurally elevated features. Steep internal structural dips severely reduce closure areas, which result in reduced potential.

Play potential

Estimates of potential for the Herschel oil play indicate a median in-place volume of 34.6 million m^3 (Fig. 18; Table 1). Predicted resource potential for gas in the Herschel play is 16.1 billion m^3 (median volume) (Fig. 19; Table 1). Predictive ranges of oil and gas resource potential vary from 2.8 to 107.9 million m^3 and 1.3 to 49.5 billion m^3, respectively. The mean number of predicted fields in both plays is 5.

The largest individual field sizes predicted for oil and gas are 15.5 million m^3 and 7.1 billion m^3, respectively (Figs. 20, 21; Table 1). No oil fields greater than 160 million m^3



Figure 17. *Schematic sketch of the trap-types in the Herschel play (from Dixon, et al., 1994).*

are expected to be present in the Herschel or Blow River highs. However, two gas fields with median estimates greater than 3 billion m^3 have been predicted.

About 81% of the play area occurs in the Yukon Territory. Under Scenario 1, where the largest undiscovered field is assumed to occur in the Yukon, the expected play potential for oil and gas is 31 million m^3 and 14.4 billion m^3, respectively (Table 2). If, however, the largest predicted field is presumed to occur either in the Northwest Territories or Alaska, the potential is reduced to 15.5 million m^3 oil and 7.2 billion m^3 gas (Table 2).



Figure 18. *Estimate of in-place oil potential of the Herschel play. Median value of probabilistic assessment is 34.6 million m^3 of in-place oil distributed in 5 fields.*



Figure 19. *Estimate of in-place gas potential of the Herschel play. Median value of probabilistic assessment is 16.1 billion m³ of in-place oil distributed in 5 fields.*



Figure 20. *Field-size-by-rank plot of the Herschel oil play. Median value of the largest predicted field size is 15.5 million m³ of in-place oil.*



Figure 21. *Field-size-by-rank plot of the Herschel gas play. Median value of the largest predicted field size is 7.1 billion m³ of in-place gas.*

YUKON COASTAL PLAIN GAS PLAY

HYDROCARBON PLAYS

Play definition

This conceptual play includes all pre-Cenomanian natural gas traps along the Yukon coastal plain extending from the northern end of the Richardson Mountains westwards to the British Mountains (Fig. 22). The play area also includes a narrow strip of inshore waters of the Beaufort Sea. It extends southward to the Barn Mountains and includes the upper reaches of the Blow River. Figure 22 shows the play may extend into Alaska to the west. However, insufficient information is available to accurately map the play boundary extensions in Alaska. Therefore, in this play assessment, all of the predictions obtained assume the play occurs exclusively in the Yukon. Gas is expected to be the principal hydrocarbon in the play.

Geology

The coastal plain play basically occupies the Rapid Depression along the Arctic coast of northern Yukon (Fig. 2). The precursory Blow Trough occupies the Rapid Depression between Cache Creek Uplift and Barn Mountains. There is a thick Mesozoic accumulation of clastic sediments in the Rapid Depression. Interbedded shale and sandstone of Jurassic to Albian age occupy the depression. These Mesozoic strata attain thicknesses of 5,000 to 10,000 m. Surrounding mountains expose Paleozoic to Proterozoic sedimentary strata. In the Barn and British mountains (Romanzof Uplift), Lower Paleozoic basinal shales, sandstone, thin-bedded carbonates and chert, Carboniferous Kayak and Kekiktuk clastic sediments and Lisburne carbonates, and Permian clastics and carbonates of the Sadlerochit Group are exposed. The thickness of the total prospect succession, including Mesozoic and Paleozoic strata, may be on the order of 5,000 to 15,000 m. Depths of this prospect succession may range from surface to 18,000 m.



Figure 22. *Yukon coastal plain gas play (from Dixon, et al., 1988). Wells discussed in the report are shown (1-Roland Bay L-41, 2-Spring River N-58, 3-Blow River E-47).*



In the Rapid Depression, Jurassic through Valanginian strata contain dominant shales of the Kingak Formation. Both the prograding clastic wedges of the sandstone-rich Bug Creek Group and the Martin Creek Formation in the Blow Trough shale-out to the west. Kamik sands are more widespread in northern Yukon except in the British Mountains to the west where an equivalent shale facies has been deposited. Late Barremian-Aptian Rat River Formation sandstones are only present on the eastern side of the Blow Trough. To the west, Rat River sands have been removed by erosion. Sediment gravity-flow deposits consisting of interbedded conglomerate, sandstone and shale of Albian age occur in the Blow Trough.

Even though the potential reservoir facies in the play seem numerous, the clastics are generally well-cemented with silica resulting in tight, low-quality reservoirs. Carbonates also exhibit low porosity characteristics but fracturing can be extensive.

Surface mapping and seismic interpretation indicate that the area is structurally complex with numerous trapping configurations occurring in association with thrust faults, normal and strike-slip faults and associated folds (Fig. 23). The major episode of deformation seems to have taken place during the Laramide orogeny, although older events have affected present trends. This implies that at least some of the structures formed prior to hydrocarbon generation and migration. Maximum burial depth when hydrocarbon generation was optimal occured during Tertiary time, so pre-existing structural traps would have captured migrating gas.

Three wells have been drilled in the play area, all in the early 1970s. All wells were dry and abandoned. Blow River E-47 in the Blow Trough stopped drilling in Albian Flysch at a depth of 4,267 m. Spring River N-58 completed drilling in Paleozoic strata at 2,136 m. The third well is designated as Roland Bay L-41 and was drilled to a total depth of 2,752 m, bottoming in Jurassic Kingak Formation. This well penetrated about 2,000 m of Jurassic and Lower Cretaceous strata of the Kingak and Mount Goodenough formations (Fig. 24, Dietrich, 1986). These dominant shale successions are unconformably overlain by the organic-rich Boundary Creek Sequence. The high vitrinite reflectance profile for all Cretaceous and Jurassic strata indicates overmature organic matter (Fig. 24). Any potential source rock, therefore, regardless of its organic type, will produce dry gas



Figure 23. *Schematic sketch of the trap-types in the Yukon coastal plain play (from Dixon, et al., 1994).*

only. Subsequent to completion of these wells, a moratorium on drilling in the area was imposed, culminating in 1987 with the establishment of a national park extending from Babbage River westward to the Alaskan border.

Many similarities are evident when comparing the Yukon coastal plain play with the Folded Ellesmerian/Pre-Mississippian play in the Arctic National Wildlife Refuge in Alaska (Dolton, *et al.*, 1987). By using the Alaskan play as an analog in this particular play assessment, some reservoir parameters and risk factors can be used.

Exploration risks

Significant marginal probabilities were assigned to presence of closure, adequate porosity, timing and preservation at the prospect level (Appendix 1). The risk for closure occurs when insufficient deformation occurs in fault-related traps for complete closure. Porosity has a high risk of being inadequate due to the well-cemented nature of many of the clastic units. Timing was considered to be a significant risk factor due to the



Figure 24. *Geological and geochemical summary of the Roland Bay L-41 well (from Dietrich, 1986).*

contemporaneous relationship of the Laramide deformational episode with a major period of hydrocarbon generation during maximum burial. Some structures were not formed before the main hydrocarbon migration episode. Uplift and deep erosion of pre-Cenomanian reservoirs during the mid-Tertiary is reflected in the high risk assigned to adequate preservation. Source and seal is most likely adequate in most prospects for gas accumulation due to the thick Jurassic and Lower Cretaceous shale succession occurring throughout the play area.

Play potential

The Yukon coastal plain play is expected to have a resource potential for gas of 8.5 billion m^3 in-place at the median value (Fig. 25; Table 1). The largest predicted field size is 5.6 billion m^3 (Fig. 26; Table 1). Three gas fields are expected in this play. Two of three fields are expected to be larger than the arbitrary 3.0 billion m^3 selected as minimum significant field size for natural gas accumulations in frontier regions of Canada. These potential volumes are applicable to the Yukon Territory only.



Figure 25. *Estimate of in-place gas potential of the Yukon coastal plain play. Median value of probabilistic assessment is 8.5 billion m^3 of in-place oil distributed in 3 fields.*

DISCUSSION OF ASSESSMENT RESULTS

RESOURCE POTENTIAL

The in-place median estimate of total oil potential for the North Coast region (including all immature and conceptual plays) is 39.8 million m^3 (250 million bbls) (Table 1; Fig. 27). (Note that the total median estimate for all oil plays is not arithmetically derived by summing the hydrocarbon potentials of individual plays. This number is derived using statistical techniques). High confidence (95% probability) and speculative (5% probability) estimates of total oil potential are 6.7 and 111 million m^3 (42 and 698 million barrels), respectively (Fig. 27). Individual field-size estimates in each play display similar probability-dependent variations. The wide range of estimates of total potential and field sizes are typical of frontier region assessments and reflect the geological uncertainties in quantifying lightly explored or conceptual exploration plays.

The median natural gas potential for all plays in the North Coast region is 38.6 billion m^3 (1.4 TCF) (Fig. 28; Table 1). The high confidence and speculative estimates for all gas



Figure 26. *Field-size-by-rank plot of the Yukon coastal plain gas play. Median value of the largest predicted field size is 6.0 billion m^3 of in-place gas.*

plays in the assessment area is 14.5 and 81.8 billion m^3 in-place (0.5 and 2.9 TCF) (Fig. 28).

Condensate occurs in a single play (South Delta–Paleozoic) where the median potential is expected to be 0.71 million m^3 or 4.5 million barrels (Table 1).

RESOURCE DISTRIBUTIONS

The greatest oil potential or volume occurs in the Herschel play (Table 1). The largest individual oil field is expected to occur in the same play, having a median size estimate of 15.5 million m^3 (97 million barrels) of in-place oil. The Herschel play also has the largest gas potential and field size in the North Coast assessment (16.1 and 7.1 billion m^3, respectively; equivalent to 0.6 and 0.25 TCF).

Taking into consideration the expected potential in the Yukon Territory alone under the two proposed scenarios (Table 2), it is clear that the Herschel play is the most significant oil and gas play with respect to both resource potential and field sizes. Under Scenario 1 where the largest predicted field is expected to occur in the Yukon Territory, 97% of the



Figure 27. *Estimate of total oil potential for the North Coast area. Median value of probabilistic assessment is 39.8 million m^3 of in-place oil distributed in 9 fields.*

oil resource and 57% of the gas resource occur in the Herschel play. When the largest field is assumed to lie outside Yukon's borders, then the proportion of gas in the Herschel play changes substantially, mostly because of the elimination of the Yukon coastal plain play in the calculation. Under Scenario 2, 98% of both the oil and gas resource occurs in the Herschel play. In any case, these calculations indicate the importance of the Herschel play in this particular assessment area. This single play, consisting of highly deformed Tertiary reservoirs and occurring disproportionately in offshore areas, infers the much greater expected oil and gas potential in the offshore region compared to onshore areas, at least in the Yukon part of the basin. No doubt, if all the other offshore plays were included as part of the assessment, this relationship would become more obvious. The relative richness of offshore plays probably reflects the proximity to the hydrocarbon kitchen adjacent to very porous and widespread Tertiary delta complexes.



Figure 28. *Estimate of total gas potential for the North Coast area. Median value of probabilistic assessment is 38.6 billion m³ of in-place oil distributed in 31 fields.*

ASSESSMENT RESULTS AND EXPLORATION HISTORY

The exploration risks calculated in this assessment suggest success rates for exploratory drilling in the North Coast area of the Yukon should average about 1 in 7. The absence of discoveries among the 3 wells drilled to date in Yukon Territory is reasonable. Historically, the first significant hydrocarbon in a frontier region is often preceded by many unsuccessful exploration wells due to incomplete knowledge of the petroleum geology in the area. This predicted success rate seems to be reasonably comparable to other frontier regions in northern Yukon (Hannigan, *et al.*, 1999; Hannigan, in press(a), (b), and (c)). The reconnaissance nature of the seismic data and the lack of wells, along with the inherent difficulty in estimating numbers of prospects associated with both structural and stratigraphic traps, make the presumption of the number of prospects quite arbitrary. Incorporating relevant future exploration data, such as improved and dense seismic survey coverage or exploratory drilling, will provide a greater degree of confidence in hydrocarbon potential estimates.

CONCLUSIONS

The hydrocarbon resource potential of the Yukon North Coast area has been evaluated through regional hydrocarbon play assessments. The quantitative assessments were derived using the Geological Survey of Canada's (PETRIMES) assessment methodology system. The Arctic coastal region of northern Yukon is underlain by a complex folded and faulted terrain: part of the Frontal morphogeological belt of the Cordilleran Fold Belt. Major tectonic and structural elements in the assessment region are the Rapid Depression, Cache Creek, Barn and Romanzof uplifts, Blow River and Herschel highs and Demarcation Subbasin. Structures such as compressional folds and strike-slip faults that were formed during the Laramide Orogeny are underlain by older tectonic features such as the Cache Creek Uplift and the Rapid Depression which originated as Mesozoic fault-bounded structures. North of the Rapid Depression are several significant large-scale tectonic features, such as the Blow River and Herschel highs and the Demarcation Subbasin.

The petroleum resource assessments include analyses of six immature and conceptual oil and gas plays, each of which incorporated the calculation or estimation of field-size parameters, numbers of prospects and exploration risks. Hydrocarbon volumes reported for these plays are total statistical estimates of the resource present 'in the ground', not the volumes that are economically producible. Individual field-size determinations are important in identifying which plays are attractive for future exploration programs.

The potential for significant hydrocarbon accumulations in the North Coast assessment region is derived by the combined presence of numerous and diverse trapping configurations, good to excellent petroleum source rocks in favourable stratal positions and reservoir-quality strata in parts of the stratigraphic succession. However, significant risks associated with lack of porosity development in Paleozoic and Mesozoic strata, and thermal maturity considerations, reduce overall hydrocarbon potential. Significant gas potential is predicted for the Herschel play in the study area. The complex geology and anticipated high exploration risks associated with all defined exploration plays in the region suggest that considerable seismic survey work and exploration drilling are required to properly evaluate the North Coast hydrocarbon potential.

The median estimate for total gas potential for all North Coast plays is 38.6 billion m^3 of in-place gas (Fig. 28; Table 1). Four fields with median sizes greater than 3 billion m^3 of in-place gas are expected in two plays, two each in the Herschel and Yukon coastal plain plays.

Total potential for oil in all North Coast plays is predicted to be 39.8 million m^3 in-place (Fig. 27; Table 1). No oil field sizes greater than 160 million m^3 are predicted to occur in the region.

REFERENCES

Alberta Energy and Utilities Board
1999: Alberta's Reserves 1998–crude oil, oil sands, gas, natural gas liquids, sulphur; Statistical Series 99-18.

Bamber, E.W. and Waterhouse, J.B.
1971: Carboniferous and Permian stratigraphy and paleontology, northern Yukon Territory, Canada; Bulletin of Canadian Petroleum Geology, v. 19, no. 1, p. 29-250.

Bird, T.D., Barclay, J.E., Campbell, R.I., and Lee, P.J.
1994: Triassic gas resources of the Western Canada Sedimentary Basin; Part I. Geological play analysis and resource assessment; Geological Survey of Canada, Bulletin 483, 66 p.

Brooks, P.W.
1986: Biological marker geochemistry of oils from the Beaufort-Mackenzie region, Arctic Canada; Bulletin of Canadian Petroleum Geology, v. 34, no. 4, p. 490-505.

Brosge, W.P., Dutro, J.T., Jr., Mangus, M.D., and Reiser, H.N.
1962: Paleozoic sequence in eastern Brooks Range, Alaska; American Association of Petroleum Geologists, Bulletin, v. 46, p. 2174-2198.

Cameron, A.R., Norris, D.K., and Pratt, K.C.
1986: Rank and other compositional data on coals and carbonaceous shale of the Kayak Formation, northern Yukon Territory, *In:* Current Research, Part B; Geological Survey of Canada, Paper 86-1B, p. 665-670.

Cecile, M.P.
1988: Corridor traverse through Barn Mountains, northernmost Yukon, *In:* Current Research, Part D; Geological Survey of Canada, Paper 88-1D, p. 99-103.

Claypool, G.E. and Magoon, L.B.
1985: Comparison of oil-source rock correlation data for Alaskan North Slope: techniques, results and conclusions, *In:* Alaska North Slope oil/rock correlation study; L.B. Magoon and G.E. Claypool (eds.), American Association of Petroleum Geologists, Studies in Geology, no. 20.

Craig, J.D., Sherwood, K.W. and Johnson, P.P.
1985: Geologic report for the Beaufort Sea planning region, Alaska; U. S. Mineral Management Service, Report MMS 85-0111, 192 p.

Davies, E.H. and Poulton, T.P.
1989: Thermal maturation data from the Jurassic rocks of northern Yukon and adjacent Northwest Territories, *In:* Current Research, Part G; Geological Survey of Canada, Paper 89-1G, p. 31-36.

Dietrich, J.R.
1986: The geology and hydrocarbon resource potential of the Beaufort Sea continental margin within the Canada–U.S.A. boundary dispute region; internal report prepared by the Petroleum Resource Secretariat, Institute of Sedimentary and Petroleum Geology; Geological Survey of Canada, 37 p.

Dietrich, J.R. and Dixon, J.
1997: Geology of the Beaufort Sea continental shelf, Chapter 12, *In:* Geology and Mineral and Hydrocarbon Potential of Northern Yukon Territory and Northwestern District of Mackenzie; D.K. Norris (ed.); Geological Survey of Canada, Bulletin 422, p. 319-322.

Dixon, J.

1986: Cretaceous to Pleistocene stratigraphy and paleogeography, northern Yukon and northwest District of Mackenzie; Bulletin of Canadian Petroleum Geology, v. 34, no. 1, p. 49-70.

1996a: Regional tectonic elements, *In:* Geological Atlas of the Beaufort-Mackenzie Area; J. Dixon (ed.); Geological Survey of Canada, Miscellaneous Report 59, p. 14-15.

1996b: Isopach map: Smoking Hills Sequence, *In:* Geological Atlas of the Beaufort-Mackenzie Area; J. Dixon (ed.); Geological Survey of Canada, Miscellaneous Report 59, p. 74.

1997: Cretaceous and Tertiary, Chapter 11, *In:* Geology and Mineral and Hydrocarbon Potential of Northern Yukon Territory and Northwestern District of Mackenzie; D.K. Norris (ed.); Geological Survey of Canada, Bulletin 422, p. 301-317.

Dixon, J. and Dietrich, J.R.

1996: Isopach map: Fish River Sequence, *In:* Geological Atlas of the Beaufort-Mackenzie Area; J. Dixon (ed.); Geological Survey of Canada, Miscellaneous Report 59, p. 75.

Dixon, J., Morrell, G.R., Dietrich, J.R., Procter, R.M., and Taylor, G.C.

1988: Petroleum resources of the Mackenzie Delta and Beaufort Sea; Geological Survey of Canada, Open File 1926, 73 p.

1994: Petroleum resources of the Mackenzie Delta and Beaufort Sea; Geological Survey of Canada, Bulletin 474, 44 p.

Dolton, G.L., Bird, K.J., and Crovelli, R.A.

1987: Assessment of in-place oil and gas resources, Chapter 22, *In:* Petroleum Geology of the Northern Part of the Arctic National Wildlife Refuge, Northeastern Alaska; K.J. Bird and L.B. Magoon (eds.); United States Geological Survey, Bulletin 1778, p. 277-298.

Dutro, J.T., Jr., Brosge, W.P. and Reiser, H.N.

1972: Significance of recently discovered Cambrian fossils and reinterpretation of Neruokpuk Formation, northeastern Alaska; American Association of Petroleum Geologists, Bulletin, v. 56, p. 808-815.

Dyke, L.D.

1973: Structural investigations in Barn Mountains, northern Yukon Territory, *In:* Report of Activities, Part A; Geological Survey of Canada, Paper 74-1A, p. 303-308.

Hamblin, A.P.

1990: Upper Paleozoic petroleum geology and potential, southern Eagle Plain, Yukon Territory; Geological Survey of Canada, Open File 2286, 49 p.

Hannigan, P.K.

in press(a): Petroleum resource assessment of the Arctic Circle/Dempster Highway Study Area; Yukon Department of Economic Development report.

in press(b): Petroleum resource assessment of the Bonnet Plume Basin; Yukon Department of Economic Development report.

in press(c): Petroleum resource assessment of the Old Crow Basin; Yukon Department of Economic Development report.

Hannigan, P.K., Dietrich, J.R., Lee, P.J., and Osadetz, K.G.

1998: Petroleum resource potential of sedimentary basins on the Pacific margin of Canada; Geological Survey of Canada, Open File 3629, 85 p.

Hannigan, P.K., Osadetz, K.G., Dixon, J. and Bird, T.

1999: Petroleum resource assessment of the Kandik basin, Yukon Territory, Canada; Yukon Department of Economic Development report, 74 p.

Hart, C.

1999: The geological framework of the Yukon Territory; Internet Web-page; www.yukonweb.com/government/geoscience/publications/summaries/framework.html.

Lane, L.S.

1988: The Rapid Fault Array: a foldbelt in Arctic Yukon, *In:* Current Research, Part A; Geological Survey of Canada, Paper 88-1A, p. 95-98.

1991: The pre-Mississippian "Neruokpuk Formation," northeastern Alaska and northwestern Yukon: review and new regional correlation; Canadian Journal of Earth Sciences, v. 28, p. 1521-1533.

1998: Latest Cretaceous-Tertiary tectonic evolution of northern Yukon and adjacent Arctic Alaska; American Association of Petroleum Geologists Bulletin, v. 82, no.7, p. 1353-1371.

Lane, L.S. and Dietrich, J.R.

1996: Structural trends and bedrock geology in the Beaufort-Mackenzie region, *In:* Geological Atlas of the Beaufort-Mackenzie Area; J. Dixon (ed.); Geological Survey of Canada, Miscellaneous Report 59, p. 10-12.

Langhus, B.G.

1980: Source and migration of hydrocarbons in the Parsons Lake area, N. W. T., Canada, *In:* Facts and Principles of World Petroleum Occurrence, A.D. Miall (ed.); Canadian Society of Petroleum Geologists, Memoir 6, p. 523-533.

Lee, P.J.

1993: Two decades of Geological Survey of Canada petroleum resource assessments; Canadian Journal of Earth Sciences, v. 30, p. 321-332.

Lee, P.J. and Tzeng, H.P.

1989: The petroleum exploration and resource evaluation system (PETRIMES): working reference guide; Institute of Sedimentary and Petroleum Geology, Calgary, Alberta, 258 p.

Lee, P.J. and Wang, P.C.C.

1990: An introduction to petroleum resource evaluation methods; Canadian Society of Petroleum Geologists, 1990 Convention on Basin Perspectives, Short Courses Program: SC-2 Petroleum Resource Evaluation, 108 p.

Link, C.M. and Bustin, R.M.

1989: Organic maturation and thermal history of Phanerozoic strata in northern Yukon and northwestern District of Mackenzie; Bulletin of Canadian Petroleum Geology, v. 37, no. 3, p. 266-292.

Link, C.M., Bustin, R.M. and Snowdon, L.R.

1989: Petroleum source potential and depositional setting of Phanerozoic strata in northern Yukon and northwestern District of Mackenzie; Bulletin of Canadian Petroleum Geology, v. 37, no. 3, p. 293-315.

Masters, C.D.

1984: Petroleum resource assessment; International Union of Geological Sciences, Publication no. 17, 157 p.

Norford, B.S.

1997: Ordovician and Silurian, Chapter 6, *In:* Geology and Mineral and Hydrocarbon Potential of Northern Yukon Territory and Northwestern District of Mackenzie; D.K. Norris (ed.); Geological Survey of Canada, Bulletin 422, p. 119-162.

Norris, D.K.

1981a: Blow River and Davidson Mountains, Yukon Territory-District of Mackenzie; Geological Survey of Canada, Map 1516A, scale 1:250,000.

1981b: Herschel Island and Demarcation Point, Yukon Territory; Geological Survey of Canada, Map 1514A, scale 1:250,000.

1997a: Geological setting, Chapter 3, *In:* Geology and Mineral and Hydrocarbon Potential of Northern Yukon Territory and Northwestern District of Mackenzie; D.K. Norris (ed.); Geological Survey of Canada, Bulletin 422, p. 21-64.

1997b: Triassic, Chapter 9, *In:* Geology and Mineral and Hydrocarbon Potential of Northern Yukon Territory and Northwestern District of Mackenzie; D.K. Norris (ed.); Geological Survey of Canada, Bulletin 422, p. 253-265.

Norris, D.K. and Dyke, L.D.

1997: Proterozoic, Chapter 4, *In:* Geology and Mineral and Hydrocarbon Potential of Northern Yukon Territory and Northwestern District of Mackenzie; D.K. Norris (ed.); Geological Survey of Canada, Bulletin 422, p. 65-83.

Northern Oil and Gas Directorate

1995a: Mackenzie Delta and Beaufort Sea, Southern Mackenzie Delta and Tuktoyaktuk Peninsula, Chapter 4, *In:* Petroleum Exploration in Northern Canada: A Guide to Oil and Gas Exploration and Potential, G.R. Morrell (ed.); Indian and Northern Affairs Canada, p. 59-64.

1995b: Mackenzie Delta and Beaufort Sea, Beaufort-Mackenzie Basin, Chapter 4, *In:* Petroleum Exploration in Northern Canada: A Guide to Oil and Gas Exploration and Potential, G.R. Morrell (ed.); Indian and Northern Affairs Canada, p. 65-73.

Podruski, J.A., Barclay, J.E., Hamblin, A.P., Lee, P.J., Osadetz, K.G., Procter, R.M., and Taylor, G.C.

1988: Conventional oil resources of Western Canada (light and medium), Part I: Resource endowment; Geological Survey of Canada, Paper 87-26, p. 1-125.

Potential Gas Committee

1990: Definitions and procedures for estimation of potential gas resources; Potential Gas Agency, Colorado School of Mines.

Poulton, T.P.

1997: Jurassic, Chapter 10, *In:* Geology and Mineral and Hydrocarbon Potential of Northern Yukon Territory and Northwestern District of Mackenzie; D.K. Norris (ed.); Geological Survey of Canada, Bulletin 422, p. 267-299.

Reed, B.L.

1968: Geology of the Lake Peters area, northeastern Brooks Range, Alaska; United States Geological Survey, Bulletin 1236.

Reinson, G.E., Lee, P.J., Warters, W., Osadetz, K.G., Bell, L.L., Price, P.R., Trollope, F., Campbell, R.I., and Barclay, J.E.

1993: Devonian gas resources of the Western Canada Sedimentary Basin; Part I: Geological play analysis and resource assessment; Geological Survey of Canada, Bulletin 452, p. 1-127.

Rice, D.D.

1986: Oil and gas assessment - methods and applications; American Association of Petroleum Geology, Studies in Geology, no. 21, 267 p.

Richards, B.C., Bamber, E.W., and Utting, J.

1997: Upper Devonian to Permian, Chapter 8, *In:* Geology and Mineral and Hydrocarbon Potential of Northern Yukon Territory and Northwestern District of Mackenzie; D.K. Norris (ed.); Geological Survey of Canada, Bulletin 422, p. 201-251.

Sable, E.G.

1977: Geology of the western Romanzof Mountains, Brooks Range, northeastern Alaska; United States Geological Survey, Professional Paper 897.

Sinclair, I.K., McAlpine, K.D., Sherwin, D.F., and McMillan, N.J.

1992: Petroleum resources of the Jeanne D'Arc Basin and environs; Part I: Geological Framework; Geological Survey of Canada, Paper 92-8.

Snowdon, L.R.

1990: Rock-eval/TOC results from 29 Beaufort-Mackenzie wells; Geological Survey of Canada, Open File Report 2192.

1996: Shell *et al.* Ulu A-35; Examples of Rock-eval/TOC analyses, *In:* Geological Atlas of the Beaufort-Mackenzie Area; J. Dixon (ed.); Geological Survey of Canada, Miscellaneous Report 59, p. 117.

Snowdon, L.R. and Powell, T.G.

1979: Families of crude oil and condensates in the Beaufort-Mackenzie Basin; Bulletin of Canadian Petroleum Geology, v. 27, p. 139-162.

Utting, J.

1989: Thermal maturity of Lower Carboniferous rocks in Yukon Territory, *In:* Current Research, Part G; Geological Survey of Canada, Paper 89-1G, p. 101-104.

Wade, J.A., Campbell, G.R., Procter, R.M., and Taylor, G.C.

1989: Petroleum resources of the Scotian Shelf; Geological Survey of Canada, Paper 88-19, 26 p.

White, D.A. and Gehman, H.M.

1979: Methods of estimating oil and gas resources; American Association of Petroleum Geologists, Bulletin, v. 63, no. 12, p. 2183-2192.

Yorath, C.J.

1991: Upper Jurassic to Paleogene assemblages, Chapter 9, *In:* Geology of the Cordilleran Orogen in Canada, H. Gabrielse and C.J. Yorath (eds.); Geological Survey of Canada, Geology of Canada, no. 4, p. 329-371.

Young, B.

1981: 3. Oil and gas, Part II, Mineral and fuel resource assessments, *In:* Assessment of Mineral and Fuel Resource Potential of the Proposed Northern Yukon National Park and Adjacent Areas; Geological Survey of Canada, Open File 760, p. 25-27.

APPENDIX 1

INPUT DATA FOR NORTH COAST HYDROCARBON ASSESSMENTS

The following tables present the probability distributions of reservoir parameters, number of prospects, and marginal probabilities of geological risk factors used as input for the various immature and conceptual statistical analyses discussed in this paper. These estimates are based on subjective opinion, partly constrained by reservoir data and information from analogous hydrocarbon-bearing basins.

1. SOUTH DELTA–MESOZOIC OIL PLAY

Table 1.1a. Probability distributions of reservoir parameters

Geological variable	Unit of measurement	Probability in upper percentiles 1.00	Probability in upper percentiles 0.50	Probability in upper percentiles 0.01	Probability in upper percentiles 0.00
Area of pool	km^2	0.11	0.7	27	34
Net pay	m	1	17	102	114
Porosity	decimal fraction	0.05	0.1	0.15	0.2
Oil saturation	decimal fraction	0.35	0.5	0.6	0.65
Oil formation volume factor	1.02	1.25	1.45	1.53	

Table 1.1b. Marginal probabilities of geological risk factors

Geological factors	Marginal probability	Play level	Prospect level
Presence of closure	0.6		x
Presence of porosity	0.50		x
Adequate seal	0.9		x
Adequate timing	0.8		x
Adequate source	0.66		x
Adequate preservation	0.7		x

Table 1.1c Probability distribution for number of prospects

Geological variable	Probability in upper percentiles 0.99	Probability in upper percentiles 0.5	Probability in upper percentiles 0.00
Number of prospects	29	35	59

2. SOUTH DELTA–MESOZOIC GAS PLAY

Table 1.2a. Probability distributions of reservoir parameters

Geological variable	Unit of measurement	Probability in upper percentiles 1.00	Probability in upper percentiles 0.50	Probability in upper percentiles 0.01	Probability in upper percentiles 0.00
Area of pool	km^2	0.11	0.7	27	34
Net pay	m	1	17	102	114
Porosity	decimal fraction	0.05	0.1	0.15	0.2
Gas saturation	decimal fraction	0.45	0.65	0.85	0.9
Gas formation volume factor	0.003	0.005	0.006	0.007	

Table 1.2b. Marginal probabilities of geological risk factors

Geological factors	Marginal probability	Play level	Prospect level
Presence of closure	0.6		x
Presence of porosity	0.50		x
Adequate seal	0.9		x
Adequate timing	0.8		x
Adequate source	0.66		x
Adequate preservation	0.7		x

Table 1.2c. Probability distribution for number of prospects

Geological variable	Probability in upper percentiles 0.99	Probability in upper percentiles 0.5	Probability in upper percentiles 0.00
Number of prospects	58	70	118

3. SOUTH DELTA–PALEOZOIC GAS PLAY

Table 1.3a. Probability distributions of reservoir parameters

Geological variable	Unit of measurement	Probability in upper percentiles 1.00	Probability in upper percentiles 0.50	Probability in upper percentiles 0.01	Probability in upper percentiles 0.00
Area of pool	km^2	0.11	0.7	27	32
Net pay	m	4	22	87	96
Porosity	decimal fraction	0.05	0.1	0.2	0.25
Gas saturation	decimal fraction	0.6	0.85	0.9	0.95
Gas formation volume factor	0.002	0.005	0.007	0.008	

Table 1.3b. Marginal probabilities of geological risk factors

Geological factors	Marginal probability	Play level	Prospect level
Presence of closure	0.6		x
Presence of porosity	0.5		x
Adequate seal	0.9		x
Adequate timing	0.8		x
Adequate source	0.8		x
Adequate preservation	0.8		x

Table 1.3c. Probability distribution for number of prospects

Geological variable	Probability in upper percentiles 0.99	Probability in upper percentiles 0.5	Probability in upper percentiles 0.00
Number of prospects	46	95	200

4. HERSCHEL OIL PLAY

Table 1.4a. Probability distributions of reservoir parameters

Geological variable	Unit of measurement	Probability in upper percentiles 1.00	Probability in upper percentiles 0.50	Probability in upper percentiles 0.01	Probability in upper percentiles 0.00
Area of pool	km^2	0.5	5	21	50
Net pay	m	3	40	200	500
Porosity	decimal fraction	0.03	0.17	0.28	0.32
Hydrocarbon saturation	decimal fraction	0.4	0.52	0.6	0.62
Oil fraction	decimal fraction	0	0.4	0.72	1
Oil formation volume factor	1.13	1.25	1.35	1.37	

Table 1.4b. Marginal probabilities of geological risk factors

Geological factors	Marginal probability	Play level	Prospect level
Presence of closure	0.7		x
Presence of porosity	0.90		x
Adequate seal	0.6		x
Adequate timing	0.8		x
Adequate source	0.7		x
Adequate preservation	0.8		x

Table 1.4c. Probability distribution for number of prospects

Geological variable	Probability in upper percentiles 0.99	Probability in upper percentiles 0.5	Probability in upper percentiles 0.00
Number of prospects	10	25	50

5. HERSCHEL GAS PLAY

Table 1.5a. Probability distributions of reservoir parameters

Geological variable	Unit of measurement	Probability in upper percentiles 1.00	Probability in upper percentiles 0.50	Probability in upper percentiles 0.01	Probability in upper percentiles 0.00
Area of pool	km^2	0.5	5	21	50
Net pay	m	3	40	200	500
Porosity	decimal fraction	0.03	0.17	0.28	0.32
Hydrocarbon saturation	decimal fraction	0.4	0.52	0.6	0.62
Gas fraction	decimal fraction	0	0.6	0.85	1
Gas formation volume factor	0.003	0.004	0.005	0.006	

Table 1.5b. Marginal probabilities of geological risk factors

Geological factors	Marginal probability	Play level	Prospect level
Presence of closure	0.7		x
Presence of porosity	0.90		x
Adequate seal	0.6		x
Adequate timing	0.8		x
Adequate source	0.7		x
Adequate preservation	0.8		x

Table 1.5c. Probability distribution for number of prospects

Geological variable	Probability in upper percentiles 0.99	Probability in upper percentiles 0.5	Probability in upper percentiles 0.00
Number of prospects	10	25	50

6. YUKON COASTAL PLAIN GAS PLAY

Table 1.6a. Probability distributions of reservoir parameters

Geological variable	Unit of measurement	Probability in upper percentiles 1.00	Probability in upper percentiles 0.50	Probability in upper percentiles 0.01	Probability in upper percentiles 0.00
Area of closure	km^2	1	9	60	185
Reservoir thickness	m	1.5	75	220	400
Trap fill	decimal fraction	0.01	0.25	0.62	1
Porosity	decimal fraction	0.03	0.1	0.17	0.22
Gas saturation	decimal fraction	0.6	0.85	0.9	0.95
Gas formation volume factor	0.002	0.005	0.007	0.008	

Table 1.6b. Marginal probabilities of geological risk factors

Geological factors	Marginal probability	Play level	Prospect level
Presence of closure	0.7		x
Presence of porosity	0.65		x
Adequate seal	0.8		x
Adequate timing	0.6		x
Adequate source	0.8		x
Adequate preservation	0.6		x

Table 1.6c. Probability distribution for number of prospects

Geological variable	Probability in upper percentiles 0.99	Probability in upper percentiles 0.5	Probability in upper percentiles 0.00
Number of prospects	5	25	40

APPENDIX 2

OUTPUT FOR NORTH COAST HYDROCARBON ASSESSMENTS

The following text presents the output generated by the PETRIMES hydrocarbon assessment program using the conceptual play analysis procedure. For each play, the PPSD, MPRO, PSRK, PSUM and PPPS modules are presented. PPSD generates the pool-size distribution using the lognormal simulation. MPRO generates the number of pools distribution and risks for the play. PSRK gives the individual pool sizes by rank and PSUM indicates the Monte Carlo simulation for the pool-ksize distribution. Immature plays with discoveries have an additional module (PPPS) which generates the play potential conditioned on the sum of discovered reserves. (Note: In text, field sizes are indicated rather than pools. In frontier conceptual plays, insufficient geological and engineering information is available to define individual pool accumulations in single structures). A PSUM module for total oil and gas potential on a basin-scale is also presented.

PETRIMES MODULE PPSD

PROSPECT OR POOL SIZE DISTRIBUTION USING LOGNORMAL OR MONTE CARLO SIMULATION

UAI C5450101
PLAY South Delta-Mesozoic oil play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date FRI, FEB 2, 2001, 9:40 AM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE ON DB?	> Y
OIL (O) OR GAS (G) ?	> O
BRITISH OR S.I. UNIT OF MEASUREMENT?	> SI
RECOVERABLE RESOURCES?	> N
DO YOU WANT TO HONOR A PERCENTILE?	> Y
UPPER PERCENTILE TO BE HONORED?	> .10000E-01

A) Basic Information

TYPE OF RESOURCE =Oil In-place
SYSTEM OF MEASUREMENT =S.I.
UNIT OF MEASUREMENT =M cu m (19)

B) Input Constant(s)

No input constant.

C) Summary

Variable		Lognormal Approximation			
IN-PLACE	mu	=-.74234	MEAN	= .95515	
POOL SIZE	sig. sq=	1.3929	S.D.	= 1.6617	
(DERIVED)	99.99% =	.59073E-02	60.00% =	.35298	15.00% = 1.6175
	99.00% =	.30564E-01	55.00% =	.41039	10.00% = 2.1601
	95.00% =	.68316E-01	50.00% =	.47600	8.00% = 2.4991
	90.00% =	.10489	45.00% =	.55210	6.00% = 2.9821
	85.00% =	.14008	40.00% =	.64189	5.00% = 3.3166
	80.00% =	.17629	35.00% =	.75008	4.00% = 3.7578
	75.00% =	.21473	30.00% =	.88388	2.00% = 5.3737
	70.00% =	.25634	25.00% =	1.0552	1.00% = 7.4131
	65.00% =	.30207	20.00% =	1.2853	.01% = 38.355

D) Pool Size Equation

NO. OF GEO-VARIABLES= 5

POOL SIZE = 1.00000 * AREA OF POOL * NET PAY * POROSITY * HYDROCARBON SATURATION / OIL FVF VOLUME FACTOR

GEOLOGICAL VARIABLES

A) Variables Used

VARIABLE	UNIT OF MEASUREMENT SYMBOL	O/G	VERSION	TYPE	NO. OF POINTS USED	UAI
AREA OF POOL	sq km	Oil	1	2	4	C5450101
NET PAY	m	Oil	1	2	4	C5450101
POROSITY	dec fr	Oil	1	2	4	C5450101
HYDROCARBON SATURATION	dec fr	Oil	1	2	4	C5450101
OIL FVF VOLUME FACTOR		Oil	1	2	4	C5450101

B) Summary

Variable		Lognormal Approximation			
AREA OF	mu	=-.35667	MEAN	= 1.2066	
POOL	sig. sq=	1.0890	S.D.	= 1.6941	
	99.99% =	.14461E-01	60.00% =	.53738	15.00% = 2.0645
	99.00% =	.61770E-01	55.00% =	.61397	10.00% = 2.6663
	95.00% =	.12579	50.00% =	.70000	8.00% = 3.0331
	90.00% =	.18378	45.00% =	.79808	6.00% = 3.5460
	85.00% =	.23734	40.00% =	.91184	5.00% = 3.8955
	80.00% =	.29085	35.00% =	1.0465	4.00% = 4.3503
	75.00% =	.34627	30.00% =	1.2099	2.00% = 5.9686
	70.00% =	.40498	25.00% =	1.4151	1.00% = 7.9326
	65.00% =	.46824	20.00% =	1.6847	.01% = 33.930
NET PAY	mu	= 2.8332	MEAN	= 19.375	
	sig. sq=	.26154	S.D.	= 10.593	
	99.99% =	2.5378	60.00% =	14.934	15.00% = 28.883
	99.00% =	5.1733	55.00% =	15.942	10.00% = 32.740
	95.00% =	7.3304	50.00% =	17.000	8.00% = 34.875
	90.00% =	8.8270	45.00% =	18.128	6.00% = 37.650
	85.00% =	10.006	40.00% =	19.352	5.00% = 39.425
	80.00% =	11.054	35.00% =	20.703	4.00% = 41.618
	75.00% =	12.040	30.00% =	22.229	2.00% = 48.595
	70.00% =	13.001	25.00% =	24.002	1.00% = 55.864
	65.00% =	13.959	20.00% =	26.144	.01% = 113.88
POROSITY	mu	=-2.3026	MEAN	= .10174	
	sig. sq=	.34487E-01	S.D.	= .19058E-01	
	99.99% =	.50125E-01	60.00% =	.95404E-01	15.00% = .12122
	99.00% =	.64920E-01	55.00% =	.97693E-01	10.00% = .12687
	95.00% =	.73678E-01	50.00% =	.10000	8.00% = .12981
	90.00% =	.78821E-01	45.00% =	.10236	6.00% = .13347
	85.00% =	.82492E-01	40.00% =	.10482	5.00% = .13572
	80.00% =	.85531E-01	35.00% =	.10742	4.00% = .13842
	75.00% =	.88227E-01	30.00% =	.11023	2.00% = .14643
	70.00% =	.90721E-01	25.00% =	.11334	1.00% = .15404
	65.00% =	.93094E-01	20.00% =	.11692	.01% = .19950

HYDROCARBON SATURATION	mu	=-.69315	MEAN	= .50124			
	sig. eq=	.49477E-02	S.D.	= .35301E-01			
	99.99%	= .38491	60.00%	= .49117	15.00%	= .53781	
	99.00%	= .42453	55.00%	= .49560	10.00%	= .54717	
	95.00%	= .44537	50.00%	= .50000	8.00%	= .55194	
	90.00%	= .45690	45.00%	= .50444	6.00%	= .55778	
	85.00%	= .46485	40.00%	= .50899	5.00%	= .56133	
	80.00%	= .47126	35.00%	= .51374	4.00%	= .56552	
	75.00%	= .47683	30.00%	= .51879	2.00%	= .57771	
	70.00%	= .48189	25.00%	= .52429	1.00%	= .58889	
	65.00%	= .48663	20.00%	= .53049	.01%	= .64950	
OIL FVN VOLUME FACTOR	mu	= .22314	MEAN	= 1.2518			
	sig. eq=	.29385E-02	S.D.	= .67910E-01			
	99.99%	= 1.0218	60.00%	= 1.2330	15.00%	= 1.3222	
	99.00%	= 1.1019	55.00%	= 1.2415	10.00%	= 1.3399	
	95.00%	= 1.1434	50.00%	= 1.2500	8.00%	= 1.3489	
	90.00%	= 1.1661	45.00%	= 1.2585	6.00%	= 1.3599	
	85.00%	= 1.1817	40.00%	= 1.2673	5.00%	= 1.3666	
	80.00%	= 1.1943	35.00%	= 1.2764	4.00%	= 1.3744	
	75.00%	= 1.2051	30.00%	= 1.2860	2.00%	= 1.3972	
	70.00%	= 1.2150	25.00%	= 1.2965	1.00%	= 1.4180	
	65.00%	= 1.2242	20.00%	= 1.3083	.01%	= 1.5292	

C) Covariance Matrix

VARIABLE					
AREA OF POOL	1.08899				
NET PAY	.00000	.26154			
POROSITY	.00000	.00000	.03449		
HYDROCARBON SATURATION	.00000	.00000	.00000	.00495	
OIL FVN VOLUME FACTOR	.00000	.00000	.00000	.00000	.00294

D) Correlation Matrix

VARIABLE					
AREA OF POOL	1.00000				
NET PAY	.00000	1.00000			
POROSITY	.00000	.00000	1.00000		
HYDROCARBON SATURATION	.00000	.00000	.00000	1.00000	
OIL FVN VOLUME FACTOR	.00000	.00000	.00000	.00000	1.00000

PETRIMES MODULE MPRO

NO. OF POOLS DISTRIBUTION AND RISKS

UAI C5450101
PLAY South Delta-Mesozoic oil play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date WED, JAN 17, 2001, 12:12 PM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE ON DB? > Y
OIL (O) OR GAS (G) ? > O

A) Risks

	GEOLOGICAL FACTOR		MARGINAL PROBABILITY
PLAY LEVEL	Overall Play Level Risk	=	1.00
PROSPECT LEVEL	Presence of Closure	(1)	.60
	Presence of Porosity	(3)	.50
	Adequate Seal	(4)	.90
	Adequate Timing	(5)	.80
	Adequate Source	(6)	.66
	Adequacy of Recovery	(9)	.70
	Overall Prospect Level Risk	=	.10
EXPLORATION RISK:		=	.10

B) No. of Prospects Distribution

Minimum = 29
Maximum = 59
Mean = 39.97
S.D. = 9.07

Frequency	No. of Prospects
99.00	29
95	30
90	31
80	32
75	32
60	34
50	35
40	40
25	47
20	50
10	55
5	57
1	59
0	59

C) No. of Pools Distribution

Minimum = 0
Maximum = 17
Mean = 3.99
S.D. = 2.10

Frequency	No. of Pools
97.88	0
95	1
90	1
80	2
75	2
60	3
50	4
40	4
25	5
20	6
10	7
5	8
1	10
0	1

Note: The no. of pools distribution is saved in the database with UDI= 62010B4

PETRIMES MODULE PSRK

INDIVIDUAL POOL SIZES BY RANK
WHERE N IS A RANDOM VARIABLE
****

UAI Č5450101
PLAY South Delta-Mesozoic oil play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date TUE, JAN 23, 2001, 12:20 PM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE ON DB ? > Y
DO YOU WANT TO USE MPRO OUTPUT? > Y
MIN. AND MAX. POOL RANKS? > 1 4
DO YOU USE LOGNORNAL ASSUMPTION? > Y
DO YOU WANT TO USE PPSD OUTPUT? > Y

A) Basic Information

TYPE OF RESOURCE =Oil In-place
SYSTEM OF MEASUREMENT =S.I.
UNIT OF MEASUREMENT =M cu m (19)

B) Lognormal Pool Size Distribution

Summary Statistics	mu =-.74234	MEAN = .95515			
	sig. sq= 1.3929	S.D. = 1.6617			
Upper Percentiles	99.99% = .59073E-02	60.00% = .35298	15.00% = 1.6175		
	99.00% = .30564E-01	55.00% = .41039	10.00% = 2.1601		
	95.00% = .68316E-01	50.00% = .47600	8.00% = 2.4991		
	90.00% = .10489	45.00% = .55210	6.00% = 2.9821		
	85.00% = .14008	40.00% = .64189	5.00% = 3.3166		
	80.00% = .17629	35.00% = .75008	4.00% = 3.7578		
	75.00% = .21473	30.00% = .88388	2.00% = 5.3737		
	70.00% = .25634	25.00% = 1.0552	1.00% = 7.4131		
	65.00% = .30207	20.00% = 1.2853	.01% = 38.355		

C) No. of Pools Distribution

Lower Support = 0
Upper Support = 17
Expectation = 3.99
Standard Deviation= 2.10

D) Pool Sizes By Rank

Pool Rank — Distribution

1	MEAN = 2.2504	S.D. = 2.8140	P(N>=r)= .97882		
	99% = .10481	75% = .78069	10% = 4.7286		
	95% = .26939	50% = 1.4679	5% = 6.6840		
	90% = .41474	25% = 2.7018	1% = 13.140		
2	MEAN = .90190	S.D. = .83927	P(N>=r)= .89821		
	99% = .50141E-01	75% = .36188	10% = 1.8576		
	95% = .12133	50% = .67353	5% = 2.4376		
	90% = .18819	25% = 1.1677	1% = 4.0505		
3	MEAN = .53469	S.D. = .46272	P(N>=r)= .74551		
	99% = .34026E-01	75% = .22023	10% = 1.0970		
	95% = .76540E-01	50% = .41050	5% = 1.4094		
	90% = .11591	25% = .70567	1% = 2.2203		
4	MEAN = .37112	S.D. = .30938	P(N>=r)= .55203		
	99% = .26794E-01	75% = .15664	10% = .75721		
	95% = .57328E-01	50% = .28836	5% = .96375		
	90% = .84734E-01	25% = .49222	1% = 1.4812		

E) The mean of the potential = 3.6164

PETRIMES MODULE PSUM

MONTE CARLO SUM SIMULATION
POOL SIZE DISTRIBUTION
****

UAI C5450101
PLAY South Delta-Mesozoic oil play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date TUE, JAN 23, 2001, 12:22 PM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE IN DATA BASE ? > Y
OIL (O) OR GAS (G) ? > O
BRITISH OR S.I. UNIT OF MEASUREMENT? > SI
RECOVERABLE RESOURCES? > N
DO YOU WANT TO USE MPRO OUTPUT? > Y
DO YOU ASSUME LOGNORMAL DISTRIBUTION? > Y
DO YOU WANT TO USE PPSD OUTPUT? > Y
DO YOU COMPUTE CONDITIONAL POTENTIAL? > N

A) Basic Information

TYPE OF RESOURCE =Oil In-place
SYSTEM OF MEASUREMENT =S.I.
UNIT OF MEASUREMENT =M cu m (19)

B) Lognormal Pool Size Distribution

Summary Statistics	mu	=-.74234	MEAN	= .95515		
	sig. sq=	1.3929	S.D.	= 1.6617		
Upper Percentiles	99.99% =	.59073E-02	60.00% =	.35298	15.00% =	1.6175
	99.00% =	.30564E-01	55.00% =	.41039	10.00% =	2.1601
	95.00% =	.68316E-01	50.00% =	.47600	8.00% =	2.4991
	90.00% =	.10489	45.00% =	.55210	6.00% =	2.9821
	85.00% =	.14008	40.00% =	.64189	5.00% =	3.3166
	80.00% =	.17629	35.00% =	.75008	4.00% =	3.7578
	75.00% =	.21473	30.00% =	.88388	2.00% =	5.3737
	70.00% =	.25634	25.00% =	1.0552	1.00% =	7.4131
	65.00% =	.30207	20.00% =	1.2853	.01% =	38.355

C) NO. OF POOLS DISTRIBUTION

Lower Support = 0
Upper Support = 17
Expectation = 3.98819
Standard Deviation= 2.09991

D) Summary Statistics for 4000 Simulations

Play Resource: (M cu m)

Minimum = .0000000E+00 Maximum = 47.66905
Expectation = 3.798451 Standard Deviation= 3.687653

EMPERICAL DISTRIBUTION:

Greater than Percentage	Play Potential
100.00	.00000E+00
95.00	.24337
90.00	.55668
85.00	.84104
80.00	1.0941
75.00	1.3875
70.00	1.6443
65.00	1.9165
60.00	2.1850
55.00	2.5113
50.00	2.8268
45.00	3.1302
40.00	3.5563
35.00	3.9955
30.00	4.4510
25.00	5.1129
20.00	5.7703
15.00	6.6875
10.00	8.1451
8.00	8.8912
6.00	9.7734
5.00	10.497
4.00	11.326
2.00	14.251
1.00	16.738
.01	46.342
.00	47.536

PETRIMES MODULE PPPS

PLAY POTENTIAL CONDITION ON
TOTAL SUM OF DISCOVERED RESERVE
****

UAI C5450101
PLAY South Delta-Mesozoic oil play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date TUE, JAN 23, 2001, 12:35 PM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE IN DATA BASE? > Y
OIL (O) OR GAS (G)? > O
BRITISH OR S.I. UNIT OF MEASUREMENT?> SI
RECOVERABLE RESOURCES? > N
CONDITION VALUE AND ITS UNIT? > 6.7000 19

A) Basic Information

TYPE OF RESOURCE =Oil In-place
SYSTEM OF MEASUREMENT =S.I.
UNIT OF MEASUREMENT =M cu m (19)

B) PLAY RESOURCE DISTRIBUTION

Summary Statistics	MEAN = 3.7985 M cu m		S.D. = 3.6877		
Upper Percentiles	100.00% = .00000E+00	55.00% = 2.5113		10.00% = 8.1451	
	95.00% = .24337	50.00% = 2.8268		8.00% = 8.8912	
	90.00% = .55668	45.00% = 3.1302		6.00% = 9.7734	
	85.00% = .84104	40.00% = 3.5563		5.00% = 10.497	
	80.00% = 1.0941	35.00% = 3.9955		4.00% = 11.326	
	75.00% = 1.3875	30.00% = 4.4510		2.00% = 14.251	
	70.00% = 1.6443	25.00% = 5.1129		1.00% = 16.738	
	65.00% = 1.9165	20.00% = 5.7703		.01% = 46.342	
	60.00% = 2.1850	15.00% = 6.6875		.00% = 47.536	

C) Summary Statistics for 5000 Simulations

Play Potential: (M cu m)

Minimum = .5783903E-03 Maximum = 40.52534
Expectation = 4.371518 Standard Deviation= 6.344550

EMPERICAL DISTRIBUTION:

Greater than Percentage	Play Potential
100.00	.57839E-03
99.00	.43201E-01
95.00	.20000
90.00	.39156
85.00	.60200
80.00	.83888
75.00	1.0402
70.00	1.2497
65.00	1.5095
60.00	1.7966
55.00	2.0800
50.00	2.3751
45.00	2.6841
40.00	2.9948
35.00	3.5396
30.00	4.1592
25.00	4.9526
20.00	6.1670
15.00	7.2779
10.00	8.8598
8.00	9.5118
6.00	13.113
5.00	17.099
4.00	21.722
2.00	30.495
1.00	34.259
.01	40.276
.00	40.500

PETRIMES MODULE PPSD

PROSPECT OR POOL SIZE DISTRIBUTION USING LOGNORMAL OR MONTE CARLO SIMULATION

****

UAI C5440101
PLAY South Delta-Mesozoic gas play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date FRI, FEB 2, 2001, 10:31 AM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE ON DB? > Y
OIL (O) OR GAS (G) ? > G
BRITISH OR S.I. UNIT OF MEASUREMENT? > SI
RECOVERABLE RESOURCES? > N
DO YOU WANT TO HONOR A PERCENTILE? > Y
UPPER PERCENTILE TO BE HONORED? > .10000E-01

A) Basic Information

TYPE OF RESOURCE =Gas In-place
SYSTEM OF MEASUREMENT =S.I.
UNIT OF MEASUREMENT =M cu m (19)

B) Input Constant(s)

No input constant.

C) Summary

Variable		Lognormal Approximation			
IN-PLACE	mu = 5.0415	MEAN = 306.47			
POOL SIZE	sig. sq= 1.3673	S.D. = 524.12			
(DERIVED)	99.99% = 1.9993	60.00% = 115.04	15.00% = 519.78		
	99.00% = 10.189	55.00% = 133.56	10.00% = 692.30		
	95.00% = 22.606	50.00% = 154.70	8.00% = 799.87		
	90.00% = 34.569	45.00% = 179.19	6.00% = 952.89		
	85.00% = 46.043	40.00% = 208.04	5.00% = 1058.7		
	80.00% = 57.822	35.00% = 242.75	4.00% = 1198.2		
	75.00% = 70.302	30.00% = 285.62	2.00% = 1707.8		
	70.00% = 83.789	25.00% = 340.42	1.00% = 2348.9		
	65.00% = 98.586	20.00% = 413.89	.01% = 11970.		

D) Pool Size Equation

NO. OF GEO-VARIABLES= 5

POOL SIZE = 1.00000 * AREA OF POOL * NET PAY * POROSITY * HYDROCARBON SATURATION / GAS FWN VOLUME FACTOR

GEOLOGICAL VARIABLES

A) Variables Used

VARIABLE	UNIT OF MEASUREMENT SYMBOL	O/G	VERSION	TYPE	NO. OF POINTS USED	UAI
AREA OF POOL	sq km	Gas	1	2	4	C5440101
NET PAY	m	Gas	1	2	4	C5440101
POROSITY	dec fr	Gas	1	2	4	C5440101
HYDROCARBON SATURATION	dec fr	Gas	1	2	4	C5440101
GAS FWN VOLUME FACTOR		Gas	1	2	4	C5440101

B) Summary

Variable		Lognormal Approximation			
AREA OF POOL	mu = -.35667	MEAN = 1.1866			
	sig. sq= 1.0555	S.D. = 1.6241			
	99.99% = .15336E-01	60.00% = .53958	15.00% = 2.0302		
	99.00% = .64138E-01	55.00% = .61522	10.00% = 2.6116		
	95.00% = .12918	50.00% = .70000	8.00% = 2.9650		
	90.00% = .18762	45.00% = .79646	6.00% = 3.4579		
	85.00% = .24135	40.00% = .90811	5.00% = 3.7932		
	80.00% = .29483	35.00% = 1.0400	4.00% = 4.2289		
	75.00% = .35007	30.00% = 1.1997	2.00% = 5.7737		
	70.00% = .40843	25.00% = 1.3997	1.00% = 7.6398		
	65.00% = .47117	20.00% = 1.6619	.01% = 31.950		
NET PAY	mu = 2.8332	MEAN = 19.375			
	sig. sq= .26154	S.D. = 10.593			
	99.99% = 2.5378	60.00% = 14.934	15.00% = 28.883		
	99.00% = 5.1733	55.00% = 15.942	10.00% = 32.740		
	95.00% = 7.3304	50.00% = 17.000	8.00% = 34.875		
	90.00% = 8.8270	45.00% = 18.128	6.00% = 37.650		
	85.00% = 10.006	40.00% = 19.352	5.00% = 39.425		
	80.00% = 11.054	35.00% = 20.703	4.00% = 41.618		
	75.00% = 12.040	30.00% = 22.229	2.00% = 48.595		
	70.00% = 13.001	25.00% = 24.002	1.00% = 55.864		
	65.00% = 13.959	20.00% = 26.144	.01% = 113.88		
POROSITY	mu = -2.3026	MEAN = .10174			
	sig. sq= .34487E-01	S.D. = .19058E-01			
	99.99% = .50125E-01	60.00% = .95404E-01	15.00% = .12122		
	99.00% = .64920E-01	55.00% = .97693E-01	10.00% = .12687		
	95.00% = .73678E-01	50.00% = .10000	8.00% = .12981		
	90.00% = .78821E-01	45.00% = .10236	6.00% = .13347		
	85.00% = .82492E-01	40.00% = .10482	5.00% = .13572		
	80.00% = .85531E-01	35.00% = .10742	4.00% = .13842		
	75.00% = .88227E-01	30.00% = .11023	2.00% = .14643		
	70.00% = .90721E-01	25.00% = .11334	1.00% = .15404		
	65.00% = .93094E-01	20.00% = .11692	.01% = .19950		

HYDROCARBON SATURATION	mu	=-.43078	MEAN	= .65248		
	sig. sq=	.76305E-02	S.D.	= .57105E-01		
	99.99% =	.46971	60.00% =	.63577	15.00% =	.71159
	99.00% =	.53047	55.00% =	.64290	10.00% =	.72699
	95.00% =	.56301	50.00% =	.65000	8.00% =	.73488
	90.00% =	.58116	45.00% =	.65717	6.00% =	.74455
	85.00% =	.59374	40.00% =	.66455	5.00% =	.75044
	80.00% =	.60393	35.00% =	.67225	4.00% =	.75741
	75.00% =	.61281	30.00% =	.68047	2.00% =	.77773
	70.00% =	.62090	25.00% =	.68945	1.00% =	.79647
	65.00% =	.62849	20.00% =	.69959	.01% =	.89950
GAS FWH VOLUME FACTOR	mu	=-5.2983	MEAN	= .50203E-02		
	sig. sq=	.81160E-02	S.D.	= .45320E-03		
	99.99% =	.35765E-02	60.00% =	.48872E-02	15.00% =	.54893E-02
	99.00% =	.40546E-02	55.00% =	.49437E-02	10.00% =	.56119E-02
	95.00% =	.43114E-02	50.00% =	.50000E-02	8.00% =	.56747E-02
	90.00% =	.44548E-02	45.00% =	.50569E-02	6.00% =	.57518E-02
	85.00% =	.45543E-02	40.00% =	.51154E-02	5.00% =	.57986E-02
	80.00% =	.46349E-02	35.00% =	.51766E-02	4.00% =	.58542E-02
	75.00% =	.47052E-02	30.00% =	.52419E-02	2.00% =	.60162E-02
	70.00% =	.47693E-02	25.00% =	.53132E-02	1.00% =	.61658E-02
	65.00% =	.48294E-02	20.00% =	.53938E-02	.01% =	.69900E-02

C) Coveriance Matrix

VARIABLE					
AREA OF POOL	1.05551				
NET PAY	.00000	.26154			
POROSITY	.00000	.00000	.03449		
HYDROCARBON SATURATION	.00000	.00000	.00000	.00763	
GAS FWH VOLUME FACTOR	.00000	.00000	.00000	.00000	.00812

D) Correlation Matrix

VARIABLE					
AREA OF POOL	1.00000				
NET PAY	.00000	1.00000			
POROSITY	.00000	.00000	1.00000		
HYDROCARBON SATURATION	.00000	.00000	.00000	1.00000	
GAS FWH VOLUME FACTOR	.00000	.00000	.00000	.00000	1.00000

PETRIMES MODULE MPRO

NO. OF POOLS DISTRIBUTION AND RISKS

UAI C5440101
PLAY South Delta-Mesozoic gas play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date MON, JAN 22, 2001, 2:08 PM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE ON DB? > Y
OIL (O) OR GAS (G) ? > G

A) Risks

	GEOLOGICAL FACTOR		MARGINAL PROBABILITY
PLAY LEVEL	Overall Play Level Risk	=	1.00
PROSPECT LEVEL	Presence of Closure	(1)	.60
	Presence of Porosity	(3)	.50
	Adequate Seal	(4)	.90
	Adequate Timing	(5)	.80
	Adequate Source	(6)	.66
	Adequate Preservation	(8)	.70
	Overall Prospect Level Risk	=	.10
EXPLORATION RISK:		=	.10

B) No. of Prospects Distribution

Minimum	= 58
Maximum	= 118
Mean	= 79.44
S.D.	= 18.14

Frequency	No. of Prospects
99.00	58
95	59
90	61
80	63
75	64
60	68
50	70
40	80
25	94
20	99
10	109
5	114
1	118
0	118

C) No. of Pools Distribution

Minimum	= 0
Maximum	= 27
Mean	= 7.93
S.D.	= 3.23

Frequency	No. of Pools
99.93	0
99	2
95	3
90	4
80	5
75	6
60	7
50	8
40	8
25	10
20	11
10	12
5	14
1	17
0	27

Note: The no. of pools distribution is saved in the database with UDI= 6201GB4

PETRIMES MODULE PSRK

INDIVIDUAL POOL SIZES BY RANK
WHERE N IS A RANDOM VARIABLE
****

UAI Cs440101
PLAY South Delta-Mesozoic gas play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date FRI, FEB 2, 2001, 10:38 AM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE ON DB ? > Y
DO YOU WANT TO USE MPRO OUTPUT? > Y
MIN. AND MAX. POOL RANKS? > 1 8
DO YOU USE LOGNORMAL ASSUMPTION? > Y
DO YOU WANT TO USE PPSD OUTPUT? > Y

A) Basic Information

TYPE OF RESOURCE =Gas In-place
SYSTEM OF MEASUREMENT =S.I.
UNIT OF MEASUREMENT =M cu m (10)

B) Lognormal Pool Size Distribution

Summary Statistics	m=	5.0415	MEAN =	306.47	
	sig. sq=	1.3673	S.D. =	524.12	
Upper Percentiles	99.99% =	1.9993	60.00% =	115.04	15.00% = 519.78
	99.00% =	10.189	55.00% =	133.56	10.00% = 692.30
	95.00% =	22.604	50.00% =	154.70	8.00% = 799.87
	90.00% =	34.569	45.00% =	179.19	6.00% = 952.89
	85.00% =	46.043	40.00% =	208.04	5.00% = 1058.7
	80.00% =	57.822	35.00% =	242.75	4.00% = 1198.2
	75.00% =	70.302	30.00% =	285.62	2.00% = 1707.8
	70.00% =	83.769	25.00% =	340.42	1.00% = 2348.9
	65.00% =	98.586	20.00% =	413.89	.01% = 11970.

C) No. of Pools Distribution

Lower Support = 0
Upper Support = 27
Expectation = 7.93
Standard Deviation= 3.23

D) Pool Sizes By Rank

Pool Rank					Distribution		
1	MEAN	= 1048.4	S.D.	= 1097.6	P(N>=r)=	.99932	
	99%	= 110.68	75%	= 452.73	10%	= 2067.1	
	95%	= 210.70	50%	= 749.83	5%	= 2828.6	
	90%	= 283.91	25%	= 1259.3	1%	= 5280.9	
2	MEAN	= 478.39	S.D.	= 357.60	P(N>=r)=	.99449	
	99%	= 50.098	75%	= 244.42	10%	= 897.91	
	95%	= 108.59	50%	= 391.46	5%	= 1135.1	
	90%	= 151.28	25%	= 608.32	1%	= 1778.3	
3	MEAN	= 300.07	S.D.	= 209.53	P(N>=r)=	.97730	
	99%	= 26.910	75%	= 155.46	10%	= 560.97	
	95%	= 63.025	50%	= 253.50	5%	= 692.82	
	90%	= 91.763	25%	= 390.33	1%	= 1025.2	
4	MEAN	= 210.49	S.D.	= 146.35	P(N>=r)=	.93669	
	99%	= 17.320	75%	= 106.51	10%	= 397.67	
	95%	= 40.378	50%	= 178.93	5%	= 487.12	
	90%	= 60.222	25%	= 278.15	1%	= 703.63	
5	MEAN	= 157.70	S.D.	= 110.85	P(N>=r)=	.86476	
	99%	= 12.772	75%	= 77.518	10%	= 301.71	
	95%	= 28.740	50%	= 133.56	5%	= 368.68	
	90%	= 42.906	25%	= 210.52	1%	= 526.65	
6	MEAN	= 124.05	S.D.	= 88.041	P(N>=r)=	.76239	
	99%	= 10.294	75%	= 59.761	10%	= 239.55	
	95%	= 22.343	50%	= 104.41	5%	= 292.62	
	90%	= 33.025	25%	= 166.47	1%	= 415.72	
7	MEAN	= 101.44	S.D.	= 72.232	P(N>=r)=	.63972	
	99%	= 8.7803	75%	= 48.483	10%	= 196.77	
	95%	= 18.517	50%	= 84.971	5%	= 240.38	
	90%	= 27.036	25%	= 136.33	1%	= 340.42	
8	MEAN	= 85.531	S.D.	= 60.723	P(N>=r)=	.51147	
	99%	= 7.7909	75%	= 40.973	10%	= 165.96	
	95%	= 16.036	50%	= 71.512	5%	= 202.70	
	90%	= 23.146	25%	= 114.86	1%	= 286.35	

E) The mean of the potential = 2353.5

PETRIMES MODULE PSUM

```
MONTE CARLO SUM SIMULATION
POOL SIZE DISTRIBUTION
**************************

UAI         CS440101
PLAY        South Delta-Mesozoic gas play
Assessor    Peter Hannigan
Geologist   Peter Hannigan
Remarks     Yukon North Slope Hydrocarbon Assessment Project
Run date    FRI, FEB  2, 2001, 10:39 AM


USER SUPPLIED PARAMETERS
------------------------
   DO YOU WANT TO STORE IN DATA BASE ?   >    Y
   OIL (O) OR GAS (G) ?                  >    G
   BRITISH OR S.I. UNIT OF MEASUREMENT?  >   SI
   RECOVERABLE RESOURCES?                >    M
   DO YOU WANT TO USE MPRO OUTPUT?       >    Y
   DO YOU ASSUME LOGNORMAL DISTRIBUTION? >    Y
   DO YOU WANT TO USE PPSD OUTPUT?       >    Y
   DO YOU COMPUTE CONDITIONAL POTENTIAL? >    N

A) Basic Information
   -----------------
   TYPE OF RESOURCE      =Gas In-place
   SYSTEM OF MEASUREMENT =S.I.
   UNIT OF MEASUREMENT   =M cu m  (19)

B) Lognormal Pool Size Distribution
   ---------------------------------
   Summary      mu     = 5.0415     MEAN   = 306.47
   Statistics   sig. sq= 1.3673     S.D.   = 524.12

   Upper        99.99% = 1.9993     60.00% = 115.04     15.00% = 519.78
   Percentiles  99.00% = 10.189     55.00% = 133.56     10.00% = 692.30
                95.00% = 22.604     50.00% = 154.70      5.00% = 799.87
                90.00% = 34.569     45.00% = 179.19      6.00% = 952.89
                85.00% = 46.043     40.00% = 208.04      5.00% = 1058.7
                80.00% = 57.822     35.00% = 242.75      4.00% = 1198.2
                75.00% = 70.302     30.00% = 285.62      2.00% = 1707.8
                70.00% = 83.789     25.00% = 340.42      1.00% = 2348.9
                65.00% = 98.586     20.00% = 413.89       .01% = 11970.

C) NO. OF POOLS DISTRIBUTION
   -------------------------------
   Lower Support     =    0
   Upper Support     =   27
   Expectation       =      7.92698
   Standard Deviation=      3.22706


D) Summary Statistics for 4000 Simulations
   ---------------------------------------

   Play Resource:    ( B cu m    )
   -------------------------------
     Minimum     = .0000000E+00Maximum          = 20.30180
     Expectation = 2.432530    Standard Deviation= 1.742724

   EMPERICAL DISTRIBUTION:
   -----------------------
```

Greater than Percentage	Play Potential
100.00	.00000E+00
99.00	.18142
95.00	.48941
90.00	.72355
85.00	.92550
80.00	1.0815
75.00	1.2620
70.00	1.4062
65.00	1.5508
60.00	1.7089
55.00	1.8698
50.00	2.0402
45.00	2.2209
40.00	2.4170
35.00	2.6301
30.00	2.8758
25.00	3.1710
20.00	3.5065
15.00	3.9387
10.00	4.5919
8.00	4.9254
6.00	5.3993
5.00	5.6884
4.00	6.0976
2.00	7.0300
1.00	8.2659
.01	18.688
.00	20.140-

PETRIMES MODULE PPSD

PROSPECT OR POOL SIZE DISTRIBUTION USING LOGNORMAL OR MONTE CARLO SIMULATION

UAI	C5460101
PLAY	South Delta-Paleozoic gas play
Assessor	Peter Hannigan
Geologist	Peter Hannigan
Remarks	Yukon North Slope Hydrocarbon Assessment Project
Run date	TUE, JAN 23, 2001, 9:42 AM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE ON DB?	> Y
OIL (O) OR GAS (G) ?	> G
BRITISH OR S.I. UNIT OF MEASUREMENT?	> SI
RECOVERABLE RESOURCES?	> N
DO YOU WANT TO HONOR A PERCENTILE?	> Y
UPPER PERCENTILE TO BE HONORED?	> .10000E-01

A) Basic Information

TYPE OF RESOURCE	=Gas In-place
SYSTEM OF MEASUREMENT	=S.I.
UNIT OF MEASUREMENT	=M cu m (19)

B) Input Constant(s)

No input constant.

C) Summary

Variable		Lognormal Approximation			
IN-PLACE	mu = 5.5676	MEAN = 498.85			
POOL SIZE	sig. sq= 1.2895	S.D. = 809.13			
(DERIVED)	99.99% = 3.8362	60.00% = 196.35		15.00% = 849.37	
	99.00% = 18.651	55.00% = 226.99		10.00% = 1122.0	
	95.00% = 40.438	50.00% = 261.80		8.00% = 1290.9	
	90.00% = 61.088	45.00% = 301.95		6.00% = 1530.1	
	85.00% = 80.694	40.00% = 349.07		5.00% = 1694.9	
	80.00% = 100.67	35.00% = 405.50		4.00% = 1911.3	
	75.00% = 121.71	30.00% = 474.88		2.00% = 2696.5	
	70.00% = 144.33	25.00% = 563.12		1.00% = 3674.8	
	65.00% = 169.02	20.00% = 680.81		.01% = 17867.	

D) Pool Size Equation

NO. OF GEO-VARIABLES= 5

POOL SIZE = 1.00000 * AREA OF POOL * NET PAY * POROSITY * HYDROCARBON SATURATION

GAS FWN VOLUME FACTOR

GEOLOGICAL VARIABLES

A) Variables Used

VARIABLE	UNIT OF MEASUREMENT SYMBOL	O/G	VERSION	TYPE	NO. OF POINTS USED	UAI
AREA OF POOL	sq km	Gas	1	2	4	C5460101
NET PAY	m	Gas	1	2	4	C5460101
POROSITY	dec fr	Gas	1	2	4	C5460101
HYDROCARBON SATURATION	dec fr	Gas	1	2	4	C5460101
GAS FWN VOLUME FACTOR		Gas	1	2	4	C5460101

B) Summary

Variable		Lognormal Approximation			
AREA OF	mu = -.35667	MEAN = 1.1866			
POOL	sig. sq= 1.0555	S.D. = 1.6241			
	99.99% = .15336E-01	60.00% = .53958		15.00% = 2.0302	
	99.00% = .64138E-01	55.00% = .61522		10.00% = 2.6116	
	95.00% = .12918	50.00% = .70000		8.00% = 2.9650	
	90.00% = .18762	45.00% = .79646		6.00% = 3.4579	
	85.00% = .24135	40.00% = .90811		5.00% = 3.7932	
	80.00% = .29483	35.00% = 1.0400		4.00% = 4.2289	
	75.00% = .35007	30.00% = 1.1997		2.00% = 5.7737	
	70.00% = .40843	25.00% = 1.3997		1.00% = 7.6398	
	65.00% = .47117	20.00% = 1.6619		.01% = 31.950	
NET PAY	mu = 3.0910	MEAN = 23.793			
	sig. sq= .15674	S.D. = 9.8014			
	99.99% = 5.0464	60.00% = 19.900		15.00% = 33.161	
	99.00% = 8.7586	55.00% = 20.932		10.00% = 36.540	
	95.00% = 11.471	50.00% = 22.000		8.00% = 38.371	
	90.00% = 13.246	45.00% = 23.122		6.00% = 40.714	
	85.00% = 14.596	40.00% = 24.321		5.00% = 42.193	
	80.00% = 15.766	35.00% = 25.626		4.00% = 43.998	
	75.00% = 16.844	30.00% = 27.076		2.00% = 49.607	
	70.00% = 17.875	25.00% = 28.734		1.00% = 55.260	
	65.00% = 18.887	20.00% = 30.699		.01% = 95.910	
POROSITY	mu = -2.3026	MEAN = .10307			
	sig. sq= .60438E-01	S.D. = .25726E-01			
	99.99% = .40080E-01	60.00% = .93962E-01		15.00% = .12902	
	99.00% = .56444E-01	55.00% = .96958E-01		10.00% = .13703	
	95.00% = .66739E-01	50.00% = .10000		8.00% = .14126	
	90.00% = .72975E-01	45.00% = .10314		6.00% = .14655	
	85.00% = .77507E-01	40.00% = .10643		5.00% = .14984	
	80.00% = .81310E-01	35.00% = .10994		4.00% = .15379	
	75.00% = .84720E-01	30.00% = .11376		2.00% = .16568	
	70.00% = .87904E-01	25.00% = .11804		1.00% = .17717	
	65.00% = .90962E-01	20.00% = .12299		.01% = .24950	

HYDROCARBON SATURATION	m:	=-.16252	MEAN	= .85038			
	sig. sq=	.88600E-03	S.D.	= .25318E-01			
	99.99% =	.76093	60.00% =	.84361	15.00% =	.87663	
	99.00% =	.79313	55.00% =	.84683	10.00% =	.88305	
	95.00% =	.80939	50.00% =	.85000	8.00% =	.88630	
	90.00% =	.81819	45.00% =	.85319	6.00% =	.89026	
	85.00% =	.82418	40.00% =	.85643	5.00% =	.89265	
	80.00% =	.82897	35.00% =	.85981	4.00% =	.89547	
	75.00% =	.83311	30.00% =	.86337	2.00% =	.90358	
	70.00% =	.83684	25.00% =	.86724	1.00% =	.91094	
	65.00% =	.84031	20.00% =	.87156	.01% =	.94950	
GAS FWH VOLUME FACTOR	m:	=-5.2983	MEAN	= .50399E-02			
	sig. sq=	.15887E-01	S.D.	= .63777E-03			
	99.99% =	.31289E-02	60.00% =	.48429E-02	15.00% =	.56978E-02	
	99.00% =	.37293E-02	55.00% =	.49214E-02	10.00% =	.58765E-02	
	95.00% =	.40638E-02	50.00% =	.50000E-02	8.00% =	.59687E-02	
	90.00% =	.42542E-02	45.00% =	.50798E-02	6.00% =	.60824E-02	
	85.00% =	.43877E-02	40.00% =	.51622E-02	5.00% =	.61519E-02	
	80.00% =	.44968E-02	35.00% =	.52488E-02	4.00% =	.62345E-02	
	75.00% =	.45925E-02	30.00% =	.53416E-02	2.00% =	.64773E-02	
	70.00% =	.46802E-02	25.00% =	.54437E-02	1.00% =	.67037E-02	
	65.00% =	.47630E-02	20.00% =	.55596E-02	.01% =	.79900E-02	

C) Covariance Matrix

VARIABLE					
AREA OF POOL	1.05551				
NET PAY	.00000	.15674			
POROSITY	.00000	.00000	.06044		
HYDROCARBON SATURATION	.00000	.00000	.00000	.00089	
GAS FWH VOLUME FACTOR	.00000	.00000	.00000	.00000	.01589

D) Correlation Matrix

VARIABLE					
AREA OF POOL	1.00000				
NET PAY	.00000	1.00000			
POROSITY	.00000	.00000	1.00000		
HYDROCARBON SATURATION	.00000	.00000	.00000	1.00000	
GAS FWH VOLUME FACTOR	.00000	.00000	.00000	.00000	1.00000

PETRIMES MODULE MPRO

NO. OF POOLS DISTRIBUTION AND RISKS

UAI C5460101
PLAY South Delta-Paleozoic gas play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date TUE, JAN 23, 2001, 9:47 AM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE ON DB? > Y
OIL (O) OR GAS (G) ? > G

A) Risks

	GEOLOGICAL FACTOR		MARGINAL PROBABILITY
PLAY LEVEL	Overall Play Level Risk	=	1.00
PROSPECT LEVEL	Presence of Closure	(1)	.60
	Presence of Porosity	(3)	.50
	Adequate Seal	(4)	.90
	Adequate Timing	(5)	.80
	Adequate Source	(6)	.80
	Adequate Preservation	(8)	.80
	Overall Prospect Level Risk	=	.14
EXPLORATION RISK:		=	.14

B) No. of Prospects Distribution

Minimum = 46
Maximum = 200
Mean = 109.25
S.D. = 45.44

Frequency	No. of Prospects
99.00	46
95	51
90	56
80	66
75	71
60	86
50	95
40	116
25	148
20	158
10	179
5	190
1	198
0	200

C) No. of Pools Distribution

Minimum = 0
Maximum = 49
Mean = 15.10
S.D. = 7.24

Frequency	No. of Pools
99.99	0
99	3
95	5
90	7
80	9
75	9
60	12
50	14
40	16
25	20
20	22
10	26
5	28
1	33
0	49

Note: The no. of pools distribution is saved in the database with UDI= 6201GB4

PETRIMES MODULE PSRK

INDIVIDUAL POOL SIZES BY RANK WHERE P IS A RANDOM VARIABLE

UAI C5460101
PLAY South Delta-Paleozoic gas play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date TUE, JAN 23, 2001, 10:37 AM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE ON DB ? > Y
DO YOU WANT TO USE MPRO OUTPUT? > Y
MIN. AND MAX. POOL RANKS? > 1 15
DO YOU USE LOGNORNAL ASSUMPTION? > Y
DO YOU WANT TO USE PPSD OUTPUT? > Y

A) Basic Information

TYPE OF RESOURCE =Gas In-place
SYSTEM OF MEASUREMENT =S.I.
UNIT OF MEASUREMENT =M cu m (19)

B) Lognormal Pool Size Distribution

Summary Statistics: mu = 5.5676, sig. sq= 1.2895, MEAN = 498.85, S.D. = 809.13

Upper Percentiles					
99.99%	= 3.8362	60.00%	= 196.35	15.00%	= 849.37
99.00%	= 18.651	55.00%	= 226.99	10.00%	= 1122.0
95.00%	= 40.438	50.00%	= 261.80	8.00%	= 1290.9
90.00%	= 61.088	45.00%	= 301.95	6.00%	= 1530.1
85.00%	= 80.694	40.00%	= 349.07	5.00%	= 1694.9
80.00%	= 100.67	35.00%	= 405.50	4.00%	= 1911.3
75.00%	= 121.71	30.00%	= 474.88	2.00%	= 2696.5
70.00%	= 144.33	25.00%	= 563.12	1.00%	= 3674.8
65.00%	= 169.02	20.00%	= 680.81	.01%	= 17867.

C) No. of Pools Distribution

Lower Support = 0
Upper Support = 49
Expectation = 15.10
Standard Deviation= 7.24

D) Pool Sizes By Rank

Pool Rank			Distribution		
1	MEAN = 2253.7	S.D. = 2029.1	P(N>=r)= .99992		
	99% = 322.85	75% = 1097.2	10% = 4254.7		
	95% = 555.12	50% = 1720.0	5% = 5652.0		
	90% = 722.72	25% = 2728.0	1% = 10012.		
2	MEAN = 1170.3	S.D. = 752.62	P(N>=r)= .99920		
	99% = 177.76	75% = 662.55	10% = 2081.9		
	95% = 328.80	50% = 1006.2	5% = 2557.3		
	90% = 434.86	25% = 1481.3	1% = 3805.1		
3	MEAN = 807.22	S.D. = 482.81	P(N>=r)= .99658		

	99%	= 105.19	75%	= 465.17	10%	= 1422.8
	95%	= 216.05	50%	= 716.43	5%	= 1704.7
	90%	= 295.23	25%	= 1042.6	1%	= 2389.8
4	MEAN	= 612.90	S.D.	= 364.58	P(N>=r)=	.98907
	99%	= 66.299	75%	= 347.15	10%	= 1088.8
	95%	= 148.54	50%	= 550.16	5%	= 1290.3
	90%	= 210.80	25%	= 805.76	1%	= 1757.9
5	MEAN	= 490.12	S.D.	= 296.26	P(N>=r)=	.97327
	99%	= 45.850	75%	= 269.28	10%	= 882.61
	95%	= 106.76	50%	= 441.23	5%	= 1040.2
	90%	= 156.41	25%	= 654.76	1%	= 1394.3
6	MEAN	= 406.18	S.D.	= 250.43	P(N>=r)=	.94640
	99%	= 34.789	75%	= 215.98	10%	= 741.58
	95%	= 81.022	50%	= 365.31	5%	= 871.17
	90%	= 120.99	25%	= 549.95	1%	= 1155.8
7	MEAN	= 346.13	S.D.	= 216.79	P(N>=r)=	.90759
	99%	= 28.411	75%	= 179.02	10%	= 638.82
	95%	= 65.097	50%	= 310.70	5%	= 748.83
	90%	= 97.891	25%	= 473.42	1%	= 986.26
8	MEAN	= 301.73	S.D.	= 190.66	P(N>=r)=	.85828
	99%	= 24.483	75%	= 153.19	10%	= 560.55
	95%	= 55.003	50%	= 270.59	5%	= 655.90
	90%	= 82.687	25%	= 415.46	1%	= 859.09
9	MEAN	= 267.91	S.D.	= 169.54	P(N>=r)=	.80152
	99%	= 21.926	75%	= 134.91	10%	= 498.79
	95%	= 48.382	50%	= 240.46	5%	= 582.66
	90%	= 72.471	25%	= 370.17	1%	= 759.73
10	MEAN	= 241.32	S.D.	= 151.98	P(N>=r)=	.74096
	99%	= 20.183	75%	= 121.69	10%	= 448.56
	95%	= 43.877	50%	= 217.17	5%	= 523.15
	90%	= 65.414	25%	= 333.65	1%	= 679.58
11	MEAN	= 219.72	S.D.	= 137.05	P(N>=r)=	.67987
	99%	= 18.949	75%	= 111.81	10%	= 406.60
	95%	= 40.693	50%	= 198.46	5%	= 473.52
	90%	= 60.370	25%	= 303.31	1%	= 613.23
12	MEAN	= 201.57	S.D.	= 124.16	P(N>=r)=	.62071
	99%	= 18.036	75%	= 104.03	10%	= 370.75
	95%	= 38.341	50%	= 182.78	5%	= 431.22
	90%	= 56.598	25%	= 277.36	1%	= 557.10
13	MEAN	= 185.82	S.D.	= 112.90	P(N>=r)=	.56500
	99%	= 17.323	75%	= 97.503	10%	= 339.50
	95%	= 36.495	50%	= 169.06	5%	= 394.50
	90%	= 53.590	25%	= 254.64	1%	= 508.78
14	MEAN	= 171.78	S.D.	= 102.99	P(N>=r)=	.51340
	99%	= 16.718	75%	= 91.630	10%	= 311.85
	95%	= 34.919	50%	= 156.64	5%	= 362.16
	90%	= 50.982	25%	= 234.35	1%	= 466.56
15	MEAN	= 159.01	S.D.	= 94.235	P(N>=r)=	.46594
	99%	= 16.153	75%	= 86.056	10%	= 287.10
	95%	= 33.443	50%	= 145.15	5%	= 333.36
	90%	= 48.521	25%	= 215.98	1%	= 429.28

E) The mean of the potential = 7203.4

PETRIMES MODULE PSUM

MONTE CARLO SUM SIMULATION
POOL SIZE DISTRIBUTION
****

UAI CS460101
PLAY South Delta-Paleozoic gas play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date TUE, JAN 23, 2001, 10:39 AM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE IN DATA BASE ? > Y
OIL (O) OR GAS (G) ? > G
BRITISH OR S.I. UNIT OF MEASUREMENT? > SI
RECOVERABLE RESOURCES? > N
DO YOU WANT TO USE MPRO OUTPUT? > Y
DO YOU ASSUME LOGNORMAL DISTRIBUTION? > Y
DO YOU WANT TO USE PPSD OUTPUT? > Y
DO YOU COMPUTE CONDITIONAL POTENTIAL? > N

A) Basic Information

TYPE OF RESOURCE =Gas In-place
SYSTEM OF MEASUREMENT =S.I.
UNIT OF MEASUREMENT =M cu m (19)

B) Lognormal Pool Size Distribution

Summary Statistics	mu	= 5.5676	MEAN	= 498.85		
	sig. sq=	1.2895	S.D.	= 809.13		
Upper Percentiles	99.99%	= 3.8362	60.00%	= 196.35	15.00%	= 849.37
	99.00%	= 18.651	55.00%	= 226.99	10.00%	= 1122.0
	95.00%	= 40.438	50.00%	= 261.80	8.00%	= 1230.9
	90.00%	= 61.088	45.00%	= 301.95	6.00%	= 1530.1
	85.00%	= 80.694	40.00%	= 349.07	5.00%	= 1694.9
	80.00%	= 100.67	35.00%	= 405.50	4.00%	= 1911.3
	75.00%	= 121.71	30.00%	= 474.88	2.00%	= 2696.5
	70.00%	= 144.33	25.00%	= 563.12	1.00%	= 3674.8
	65.00%	= 169.02	20.00%	= 680.81	.01%	= 17867.

C) NO. OF POOLS DISTRIBUTION

Lower Support = 0
Upper Support = 49
Expectation = 15.10272
Standard Deviation= 7.24418

D) Summary Statistics for 4000 Simulations

Play Resource: (B cu m)

Minimum = .1544554 Maximum = 33.45845
Expectation = 7.523363 Standard Deviation= 4.663376

EMPERICAL DISTRIBUTION:

Greater than Percentage	Play Potential
100.00	.15446
99.00	.87468
95.00	1.6994
90.00	2.3555
85.00	2.9483
80.00	3.4788
75.00	3.9972
70.00	4.5850
65.00	5.0365
60.00	5.5197
55.00	6.0653
50.00	6.6614
45.00	7.2674
40.00	7.8857
35.00	8.5103
30.00	9.2216
25.00	10.064
20.00	11.098
15.00	12.215
10.00	13.715
8.00	14.470
6.00	15.512
5.00	16.536
4.00	17.344
2.00	19.535
1.00	21.752
.01	33.164
.00	33.429

PETRIMES MODULE PPPS

PLAY POTENTIAL CONDITION ON
TOTAL SUM OF DISCOVERED RESERVE
****

URI C5460101
PLAY South Delta-Paleozoic gas play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date TUE, JAN 23, 2001, 10:41 AM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE IN DATA BASE? > Y
OIL (O) OR GAS (G)? > G
BRITISH OR S.I. UNIT OF MEASUREMENT?> SI
RECOVERABLE RESOURCES? > M
CONDITION VALUE AND ITS UNIT? > 311.00 19

A) Basic Information

TYPE OF RESOURCE =Gas In-place
SYSTEM OF MEASUREMENT =S.I.
UNIT OF MEASUREMENT =M cu m (19)

B) PLAY RESOURCE DISTRIBUTION

Summary Statistics: MEAN = 7.5234 S.D. = 4.6634 B cu m

Upper Percentiles	100.00% = .15446	55.00% = 6.0653	8.00% = 14.470
	99.00% = .87468	50.00% = 6.6614	6.00% = 15.512
	95.00% = 1.6994	45.00% = 7.2674	5.00% = 16.536
	90.00% = 2.3555	40.00% = 7.8857	4.00% = 17.344
	85.00% = 2.9483	35.00% = 8.5103	2.00% = 19.535
	80.00% = 3.4788	30.00% = 9.2216	1.00% = 21.752
	75.00% = 3.9972	25.00% = 10.064	.01% = 33.164
	70.00% = 4.5850	20.00% = 11.098	.00% = 33.429
	65.00% = 5.0365	15.00% = 12.215	
	60.00% = 5.5197	10.00% = 13.715	

C) Summary Statistics for 5000 Simulations

```
--------**--------------------------------

Play Potential:   ( B cu m    )
-------------------------------
  Minimum      = .1047840E-01Maximum          = 32.41308
  Expectation = 7.279167    Standard Deviation= 4.818757

EMPERICAL DISTRIBUTION:
-----------------------

  Greater than    Play
  Percentage      Potential
  ------------    -----------
     100.00         .10478E-01
      99.00         .60054
      95.00         1.3824
      90.00         2.0107
      85.00         2.5747
      80.00         3.1288
      75.00         3.6557
      70.00         4.2706
      65.00         4.7228
      60.00         5.1871
      55.00         5.8006
      50.00         6.3701
      45.00         7.0142
      40.00         7.6513
      35.00         8.2200
      30.00         8.9133
      25.00         9.8504
      20.00         10.863
      15.00         12.036
      10.00         13.482
       8.00         14.405
       6.00         15.598
       5.00         16.573
       4.00         17.544
       2.00         19.687
       1.00         21.719
        .01         32.318
        .00         32.404
```

PETRIMES MODULE PPSD

```
PROSPECT OR POOL SIZE DISTRIBUTION USING
LOGNORMAL OR MONTE CARLO SIMULATION
*******************************************

UAI         CS4E0101
PLAY        Herschel oil play
Assessor    Peter Hannigan
Geologist   Peter Hannigan
Remarks     Yukon North Slope Hydrocarbon Assessment Project
Run date    FRI, JAN 26, 2001, 12:08 PM


USER SUPPLIED PARAMETERS
------------------------
   DO YOU WANT TO STORE ON DB?           >    Y
   OIL (O) OR GAS (G) ?                  >    O
   BRITISH OR S.I. UNIT OF MEASUREMENT?  >   SI
   RECOVERABLE RESOURCES?                >    N
   DO YOU WANT TO HONOR A PERCENTILE?    > Y
   UPPER PERCENTILE TO BE HONORED?       > .10000E-01

A) Basic Information
   -----------------
   TYPE OF RESOURCE      =Oil In-place
   SYSTEM OF MEASUREMENT =S.I.
   UNIT OF MEASUREMENT   =M cu m  (19)

B) Input Constant(s)
   -----------------

   No input constant.

C) Summary
   -------

   Variable                          Lognormal Approximation

   IN-PLACE      mu     = 1.7330     MEAN   = 9.0178
   POOL SIZE     sig. sq= .93241     S.D.   = 11.193
   (DERIVED)     99.99% = .15596     60.00% = 4.4298     15.00% = 15.391
                 99.00% = .59848     55.00% = 5.0111     10.00% = 19.501
                 95.00% = 1.1557     50.00% = 5.6576      8.00% = 21.972
                 90.00% = 1.6413     45.00% = 6.3875      6.00% = 25.389
                 85.00% = 2.0797     40.00% = 7.2256      5.00% = 27.696
                 80.00% = 2.5101     35.00% = 8.2077      4.00% = 30.676
                 75.00% = 2.9497     30.00% = 9.3874      2.00% = 41.105
                 70.00% = 3.4097     25.00% = 10.851      1.00% = 53.482
                 65.00% = 3.8998     20.00% = 12.752       .01% = 205.23

D) Pool Size Equation
   ------------------

   NO. OF GEO-VARIABLES= 4

   POOL SIZE =        1.00000
              * AREA OF      * NET PAY      * POROSITY      * HYDROCARBON
                POOL                                        SATURATION
              * OIL FRACTION
                OIL FVF
                VOLUME
                FACTOR
```

GEOLOGICAL VARIABLES

A) Variables Used

VARIABLE	UNIT OF MEASUREMENT SYMBOL	O/G	VERSION	TYPE	NO. OF POINTS USED	UAI
AREA OF POOL	sq km	Oil	1	2	4	C54E0101
NET PAY	m	Oil	1	2	4	C54E0101
POROSITY	dec fr	Oil	1	2	4	C54E0101
HYDROCARBON SATURATION	dec fr	Oil	1	2	4	C54E0101
OIL FRACTION	dec fr	Oil	1	2	4	C54E0101
OIL FWN VOLUME FACTOR		Oil	1	2	4	C54E0101

B) Summary

Variable		Lognormal Approximation		
AREA OF POOL	mu = 1.6094	MEAN = 6.0505		
	sig. sq= .38140	S.D. = 4.1229		
	99.99% = .50292	60.00% = 4.2758	15.00% = 9.4831	
	99.00% = 1.1886	55.00% = 4.6266	10.00% = 11.033	
	95.00% = 1.8105	50.00% = 5.0000	8.00% = 11.908	
	90.00% = 2.2659	45.00% = 5.4035	6.00% = 13.061	
	85.00% = 2.6363	40.00% = 5.8468	5.00% = 13.808	
	80.00% = 2.9733	35.00% = 6.3433	4.00% = 14.741	
	75.00% = 3.2966	30.00% = 6.9122	2.00% = 17.775	
	70.00% = 3.6168	25.00% = 7.5836	1.00% = 21.034	
	65.00% = 3.9412	20.00% = 8.4081	.01% = 49.710	
NET PAY	mu = 3.6889	MEAN = 50.320		
	sig. sq= .45903	S.D. = 38.406		
	99.99% = 3.2193	60.00% = 33.691	15.00% = 80.728	
	99.00% = 8.2708	55.00% = 36.735	10.00% = 95.312	
	95.00% = 13.124	50.00% = 40.000	8.00% = 103.63	
	90.00% = 16.787	45.00% = 43.555	6.00% = 114.69	
	85.00% = 19.820	40.00% = 47.490	5.00% = 121.91	
	80.00% = 22.616	35.00% = 51.932	4.00% = 130.97	
	75.00% = 25.328	30.00% = 57.064	2.00% = 160.83	
	70.00% = 28.039	25.00% = 63.172	1.00% = 193.45	
	65.00% = 30.809	20.00% = 70.746	.01% = 497.00	
POROSITY	mu = -1.7720	MEAN = .17247		
	sig. sq= .28812E-01	S.D. = .29487E-01		
	99.99% = .90426E-01	60.00% = .16284	15.00% = .20270	
	99.00% = .11454	55.00% = .16641	10.00% = .21131	
	95.00% = .12859	50.00% = .17000	8.00% = .21579	
	90.00% = .13677	45.00% = .17367	6.00% = .22134	
	85.00% = .14258	40.00% = .17747	5.00% = .22475	
	80.00% = .14737	35.00% = .18149	4.00% = .22883	
	75.00% = .15161	30.00% = .18583	2.00% = .24091	
	70.00% = .15552	25.00% = .19062	1.00% = .25231	
	65.00% = .15924	20.00% = .19611	.01% = .31960	
HYDROCARBON SATURATION	mu = -.65393	MEAN = .52058		
	sig. sq= .22286E-02	S.D. = .24589E-01		
	99.99% = .43627	60.00% = .51382	15.00% = .54608	
	99.00% = .46592	55.00% = .51692	10.00% = .55243	
	95.00% = .48115	50.00% = .52000	8.00% = .55566	
	90.00% = .48947	45.00% = .52309	6.00% = .55960	
	85.00% = .49517	40.00% = .52626	5.00% = .56199	
	80.00% = .49974	35.00% = .52955	4.00% = .56480	
	75.00% = .50370	30.00% = .53303	2.00% = .57294	
	70.00% = .50728	25.00% = .53682	1.00% = .58036	
	65.00% = .51063	20.00% = .54108	.01% = .61980	
OIL FRACTION	mu = -.91629	MEAN = .41225		
	sig. sq= .60332E-01	S.D. = .10281		
	99.99% = .16045	60.00% = .37587	15.00% = .51597	
	99.00% = .22589	55.00% = .38784	10.00% = .54798	
	95.00% = .26705	50.00% = .40000	8.00% = .56486	
	90.00% = .29198	45.00% = .41254	6.00% = .58602	
	85.00% = .31010	40.00% = .42568	5.00% = .59913	
	80.00% = .32530	35.00% = .43971	4.00% = .61491	
	75.00% = .33893	30.00% = .45499	2.00% = .66243	
	70.00% = .35166	25.00% = .47207	1.00% = .70831	
	65.00% = .36388	20.00% = .49186	.01% = .99720	
OIL FWN VOLUME FACTOR	mu = .22314	MEAN = 1.2504		
	sig. sq= .60560E-03	S.D. = .30775E-01		
	99.99% = 1.1407	60.00% = 1.2422	15.00% = 1.2823	
	99.00% = 1.1804	55.00% = 1.2461	10.00% = 1.2901	
	95.00% = 1.2004	50.00% = 1.2500	8.00% = 1.2940	
	90.00% = 1.2112	45.00% = 1.2539	6.00% = 1.2988	
	85.00% = 1.2185	40.00% = 1.2578	5.00% = 1.3016	
	80.00% = 1.2244	35.00% = 1.2619	4.00% = 1.3050	
	75.00% = 1.2294	30.00% = 1.2662	2.00% = 1.3148	
	70.00% = 1.2340	25.00% = 1.2709	1.00% = 1.3236	
	65.00% = 1.2382	20.00% = 1.2762	.01% = 1.3698	

C) Covariance Matrix

VARIABLE					
AREA OF POOL	.38140				
NET PAY	.00000	.45903			
POROSITY	.00000	.00000	.02881		
HYDROCARBON SATURATION	.00000	.00000	.00000	.00223	
OIL FRACTION	.00000	.00000	.00000	.00000	.06033
OIL FWN VOLUME FACTOR	.00000	.00000	.00000	.00000	.00000

VARIABLE	
OIL FWN VOLUME FACTOR	.00061

D) Correlation Matrix

VARIABLE					
AREA OF POOL	1.00000				
NET PAY	.00000	1.00000			
POROSITY	.00000	.00000	1.00000		
HYDROCARBON SATURATION	.00000	.00000	.00000	1.00000	
OIL FRACTION	.00000	.00000	.00000	.00000	1.00000
OIL FVF VOLUME FACTOR	.00000	.00000	.00000	.00000	.00000

VARIABLE	
OIL FVF VOLUME FACTOR	1.00000

PETRIMES MODULE MPRO

NO. OF POOLS DISTRIBUTION AND RISKS

UAI C54E0101
PLAY Herschel oil play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date FRI, JAN 26, 2001, 12:09 PM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE ON DB? > Y
OIL (O) OR GAS (G) ? > O

A) Risks

	GEOLOGICAL FACTOR		MARGINAL PROBABILITY
PLAY LEVEL	Overall Play Level Risk	=	1.00
PROSPECT LEVEL	Presence of Closure	(1)	.70
	Presence of Porosity	(3)	.90
	Adequate Seal	(4)	.60
	Adequate Timing	(5)	.80
	Adequate Source	(6)	.70
	Adequate Preservation	(8)	.80
	Overall Prospect Level Risk	=	.17
EXPLORATION RISK:		=	.17

B) No. of Prospects Distribution

Minimum	= 10
Maximum	= 50
Mean	= 27.92
S.D.	= 11.72

Frequency	No. of Prospects
99.00	10
95	12
90	13
80	16
75	18
60	22
50	25
40	30
25	38
20	40
10	45
5	48
1	50
0	50

C) No. of Pools Distribution

Minimum	= 0
Maximum	= 21
Mean	= 4.73
S.D.	= 2.80

Frequency	No. of Pools
97.40	0
95	1
90	1
80	2
75	3
60	4
50	4
40	5
25	6
20	7
10	9
5	10
1	12
0	21

Note: The no. of pools distribution is saved in the database with UDI=6201OB4

PETRIMES MODULE PSRK

INDIVIDUAL POOL SIZES BY RANK
WHERE N IS A RANDOM VARIABLE
****

UAI CS4E0101
PLAY Herschel oil play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date FRI, JAN 26, 2001, 12:10 PM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE ON DB ? > Y
DO YOU WANT TO USE MPRO OUTPUT? > Y
MIN. AND MAX. POOL RANKS? > 1 5
DO YOU USE LOGNORMAL ASSUMPTION? > Y
DO YOU WANT TO USE PPSD OUTPUT? > Y

A) Basic Information

TYPE OF RESOURCE =Oil In-place
SYSTEM OF MEASUREMENT =S.I.
UNIT OF MEASUREMENT =M cu m (19)

B) Lognormal Pool Size Distribution

Summary Statistics	mu	= 1.7330	MEAN	= 9.0178	
	sig. sq	= .93241	S.D.	= 11.193	
Upper Percentiles	99.99%	= .15596	60.00%	= 4.4298	15.00% = 15.391
	99.00%	= .59848	55.00%	= 5.0111	10.00% = 19.501
	95.00%	= 1.1557	50.00%	= 5.6576	8.00% = 21.972
	90.00%	= 1.6413	45.00%	= 6.3875	6.00% = 25.389
	85.00%	= 2.0797	40.00%	= 7.2256	5.00% = 27.696
	80.00%	= 2.5101	35.00%	= 8.2077	4.00% = 30.676
	75.00%	= 2.9497	30.00%	= 9.3874	2.00% = 41.105
	70.00%	= 3.4097	25.00%	= 10.851	1.00% = 53.482
	65.00%	= 3.8998	20.00%	= 12.752	.01% = 205.23

C) No. of Pools Distribution

Lower Support = 0
Upper Support = 21
Expectation = 4.73
Standard Deviation = 2.80

D) Pool Sizes By Rank

Pool Rank					Distribution	
1	MEAN	= 20.290	S.D.	= 18.379	P(N>=r)	= .97399
	99%	= 1.6600	75%	= 9.2188	10%	= 39.684
	95%	= 3.7164	50%	= 15.499	5%	= 52.367
	90%	= 5.3906	25%	= 25.345	1%	= 90.150
2	MEAN	= 10.240	S.D.	= 7.3619	P(N>=r)	= .89343
	99%	= .96181	75%	= 5.1390	10%	= 19.342
	95%	= 2.0305	50%	= 8.5941	5%	= 23.977
	90%	= 2.9513	25%	= 13.382	1%	= 35.835
3	MEAN	= 6.9831	S.D.	= 4.7122	P(N>=r)	= .76446
	99%	= .72994	75%	= 3.5827	10%	= 13.058
	95%	= 1.4571	50%	= 5.9920	5%	= 15.878
	90%	= 2.0780	25%	= 9.2387	1%	= 22.637
4	MEAN	= 5.3403	S.D.	= 3.4743	P(N>=r)	= .61665
	99%	= .61565	75%	= 2.7948	10%	= 9.8998
	95%	= 1.1827	50%	= 4.6277	5%	= 11.927
	90%	= 1.6555	25%	= 7.0817	1%	= 16.619
5	MEAN	= 4.3322	S.D.	= 2.7386	P(N>=r)	= .47506
	99%	= .54402	75%	= 2.3146	10%	= 7.9583
	95%	= 1.0163	50%	= 3.7785	5%	= 9.5351
	90%	= 1.4008	25%	= 5.7350	1%	= 13.111

E) The mean of the potential = 39.601

PETRIMES MODULE PSUM

MONTE CARLO SUM SIMULATION
POOL SIZE DISTRIBUTION
****

UAI Cs4E0101
PLAY Herschel oil play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date FRI, JAN 26, 2001, 12:13 PM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE IN DATA BASE ?	> Y
OIL (O) OR GAS (G) ?	> O
BRITISH OR S.I. UNIT OF MEASUREMENT?	> SI
RECOVERABLE RESOURCES?	> N
DO YOU WANT TO USE MPRO OUTPUT?	> Y
DO YOU ASSUME LOGNORMAL DISTRIBUTION?	> Y
DO YOU WANT TO USE PPSD OUTPUT?	> Y
DO YOU COMPUTE CONDITIONAL POTENTIAL?	> N

A) Basic Information

TYPE OF RESOURCE =Oil In-place
SYSTEM OF MEASUREMENT =S.I.
UNIT OF MEASUREMENT =M cu m (19)

B) Lognormal Pool Size Distribution

Summary Statistics	mu	= 1.7330	MEAN	= 9.0178	
	sig. sq	= .93241	S.D.	= 11.193	
Upper Percentiles	99.99%	= .15596	60.00%	= 4.4298	15.00% = 15.391
	99.00%	= .59848	55.00%	= 5.0111	10.00% = 19.501
	95.00%	= 1.1557	50.00%	= 5.6576	8.00% = 21.972
	90.00%	= 1.6413	45.00%	= 6.3875	6.00% = 25.389
	85.00%	= 2.0797	40.00%	= 7.2256	5.00% = 27.696
	80.00%	= 2.5101	35.00%	= 8.2077	4.00% = 30.676
	75.00%	= 2.9497	30.00%	= 9.3874	2.00% = 41.105
	70.00%	= 3.4097	25.00%	= 10.851	1.00% = 53.482
	65.00%	= 3.8998	20.00%	= 12.752	.01% = 205.23

C) NO. OF POOLS DISTRIBUTION

Lower Support	= 0
Upper Support	= 21
Expectation	= 4.72808
Standard Deviation	= 2.80484

D) Summary Statistics for 4000 Simulations

Play Resource: (M cu m)

Minimum = .0000000E+00 Maximum = 349.0305
Expectation = 42.40630 Standard Deviation = 34.55067

EMPERICAL DISTRIBUTION:

Greater than Percentage	Play Potential
100.00	.00000E+00
95.00	2.7805
90.00	6.8992
85.00	10.490
80.00	13.667
75.00	16.913
70.00	20.214
65.00	23.473
60.00	27.095
55.00	30.753
50.00	34.605
45.00	38.698
40.00	43.242
35.00	48.156
30.00	52.590
25.00	59.358
20.00	65.791
15.00	75.218
10.00	88.821
8.00	95.161
6.00	101.91
5.00	107.90
4.00	114.67
2.00	132.80
1.00	156.25
.01	324.33
.00	346.56

PETRIMES MODULE PPSD

PROSPECT OR POOL SIZE DISTRIBUTION USING LOGNORMAL OR MONTE CARLO SIMULATION

UAI C54D0101
PLAY Herschel gas play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date FRI, JAN 26, 2001, 11:13 AM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE ON DB? > Y
OIL (O) OR GAS (G) ? > G
BRITISH OR S.I. UNIT OF MEASUREMENT? > SI
RECOVERABLE RESOURCES? > N
DO YOU WANT TO HONOR A PERCENTILE? > Y
UPPER PERCENTILE TO BE HONORED? > .10000E-01

A) Basic Information

TYPE OF RESOURCE =Gas In-place
SYSTEM OF MEASUREMENT =S.I.
UNIT OF MEASUREMENT =M cu m (19)

B) Input Constant(s)

No input constant.

C) Summary

Variable			Lognormal Approximation		
IN-PLACE	mu	= 7.8831	MEAN = 4163.4		
POOL SIZE	sig. sq=	.90202	S.D. = 5038.5		
(DERIVED)	99.99%	= 77.552	60.00% = 2084.9	15.00%	= 7096.9
	99.00%	= 291.09	55.00% = 2353.7	10.00%	= 8957.2
	95.00%	= 556.06	50.00% = 2652.0	8.00%	= 10072.
	90.00%	= 785.19	45.00% = 2988.2	6.00%	= 11611.
	85.00%	= 991.01	40.00% = 3373.4	5.00%	= 12648.
	80.00%	= 1192.4	35.00% = 3823.9	4.00%	= 13985.
	75.00%	= 1397.5	30.00% = 4363.9	2.00%	= 18650.
	70.00%	= 1611.7	25.00% = 5032.5	1.00%	= 24161.
	65.00%	= 1839.2	20.00% = 5898.2	.01%	= 90689.

D) Pool Size Equation

NO. OF GEO-VARIABLES= 6

POOL SIZE = 1.00000 * AREA OF POOL * NET PAY * POROSITY * HYDROCARBON SATURATION * GAS FRACTION

GAS FWN VOLUME FACTOR

GEOLOGICAL VARIABLES

A) Variables Used

VARIABLE	UNIT OF MEASUREMENT SYMBOL	O/G	VERSION	TYPE	NO. OF POINTS USED	UAI
AREA OF POOL	sq km	Gas	1	2	4	C54D0101
NET PAY	m	Gas	1	2	4	C54D0101
POROSITY	dec fr	Gas	1	2	4	C54D0101
HYDROCARBON SATURATION	dec fr	Gas	1	2	4	C54D0101
GAS FRACTION	dec fr	Gas	1	2	4	C54D0101
GAS FWN VOLUME FACTOR		Gas	1	2	4	C54D0101

B) Summary

Variable			Lognormal Approximation		
AREA OF	mu	= 1.6094	MEAN = 6.0505		
POOL	sig. sq=	.38140	S.D. = 4.1229		
	99.99%	= .50292	60.00% = 4.2758	15.00%	= 9.4831
	99.00%	= 1.1886	55.00% = 4.6266	10.00%	= 11.033
	95.00%	= 1.8105	50.00% = 5.0000	8.00%	= 11.908
	90.00%	= 2.2659	45.00% = 5.4035	6.00%	= 13.061
	85.00%	= 2.6363	40.00% = 5.8468	5.00%	= 13.808
	80.00%	= 2.9733	35.00% = 6.3433	4.00%	= 14.741
	75.00%	= 3.2966	30.00% = 6.9122	2.00%	= 17.775
	70.00%	= 3.6168	25.00% = 7.5836	1.00%	= 21.034
	65.00%	= 3.9412	20.00% = 8.4081	.01%	= 49.710
NET PAY	mu	= 3.6889	MEAN = 50.320		
	sig. sq=	.45903	S.D. = 38.406		
	99.99%	= 3.2193	60.00% = 33.691	15.00%	= 80.728
	99.00%	= 8.2708	55.00% = 36.735	10.00%	= 95.312
	95.00%	= 13.124	50.00% = 40.000	8.00%	= 103.63
	90.00%	= 16.787	45.00% = 43.555	6.00%	= 114.69

	85.00% = 19.820	40.00% = 47.490	5.00% = 121.91		
	80.00% = 22.616	35.00% = 51.932	4.00% = 130.97		
	75.00% = 25.328	30.00% = 57.064	2.00% = 160.83		
	70.00% = 28.039	25.00% = 63.172	1.00% = 193.45		
	65.00% = 30.809	20.00% = 70.746	.01% = 497.00		
POROSITY	mu =-1.7720	MEAN = .17247			
	sig. sq= .28812E-01	S.D. = .29487E-01			
	99.99% = .90426E-01	60.00% = .16284	15.00% = .20270		
	99.00% = .11454	55.00% = .16641	10.00% = .21131		
	95.00% = .12859	50.00% = .17000	8.00% = .21579		
	90.00% = .13677	45.00% = .17367	6.00% = .22134		
	85.00% = .14258	40.00% = .17747	5.00% = .22475		
	80.00% = .14737	35.00% = .18149	4.00% = .22883		
	75.00% = .15161	30.00% = .18583	2.00% = .24091		
	70.00% = .15552	25.00% = .19062	1.00% = .25231		
	65.00% = .15924	20.00% = .19611	.01% = .31960		
HYDROCARBON SATURATION	mu =-.65393	MEAN = .52058			
	sig. sq= .22286E-02	S.D. = .24589E-01			
	99.99% = .43627	60.00% = .51382	15.00% = .54608		
	99.00% = .46592	55.00% = .51692	10.00% = .55243		
	95.00% = .48115	50.00% = .52000	8.00% = .55566		
	90.00% = .48947	45.00% = .52309	6.00% = .55960		
	85.00% = .49517	40.00% = .52626	5.00% = .56199		
	80.00% = .49974	35.00% = .52955	4.00% = .56480		
	75.00% = .50370	30.00% = .53303	2.00% = .57294		
	70.00% = .50728	25.00% = .53682	1.00% = .58036		
	65.00% = .51063	20.00% = .54108	.01% = .61980		
GAS FRACTION	mu =-.51083	MEAN = .60565			
	sig. sq= .18756E-01	S.D. = .83335E-01			
	99.99% = .36054	60.00% = .57954	15.00% = .69151		
	99.00% = .43630	55.00% = .58976	10.00% = .71511		
	95.00% = .47898	50.00% = .60000	8.00% = .72731		
	90.00% = .50342	45.00% = .61042	6.00% = .74238		
	85.00% = .52060	40.00% = .62118	5.00% = .75159		
	80.00% = .53468	35.00% = .63251	4.00% = .76257		
	75.00% = .54706	30.00% = .64468	2.00% = .79488		
	70.00% = .55842	25.00% = .65806	1.00% = .82512		
	65.00% = .56916	20.00% = .67330	.01% = .99850		
GAS FWN VOLUME FACTOR	mu =-5.5215	MEAN = .40236E-02			
	sig. sq= .11789E-01	S.D. = .43816E-03			
	99.99% = .26711E-02	60.00% = .38915E-02	15.00% = .44764E-02		
	99.00% = .31072E-02	55.00% = .39458E-02	10.00% = .45972E-02		
	95.00% = .33458E-02	50.00% = .40000E-02	8.00% = .46592E-02		
	90.00% = .34804E-02	45.00% = .40549E-02	6.00% = .47356E-02		
	85.00% = .35743E-02	40.00% = .41116E-02	5.00% = .47821E-02		
	80.00% = .36507E-02	35.00% = .41709E-02	4.00% = .48374E-02		
	75.00% = .37175E-02	30.00% = .42344E-02	2.00% = .49992E-02		
	70.00% = .37786E-02	25.00% = .43039E-02	1.00% = .51494E-02		
	65.00% = .38361E-02	20.00% = .43827E-02	.01% = .59900E-02		

C) Covariance Matrix

VARIABLE					
AREA OF POOL	.38140				
NET PAY	.00000	.45903			
POROSITY	.00000	.00000	.02881		
HYDROCARBON SATURATION	.00000	.00000	.00000	.00223	
GAS FRACTION	.00000	.00000	.00000	.00000	.01876
GAS FWN VOLUME FACTOR	.00000	.00000	.00000	.00000	.00000

VARIABLE	
GAS FWN VOLUME FACTOR	.01179

D) Correlation Matrix

VARIABLE					
AREA OF POOL	1.00000				
NET PAY	.00000	1.00000			
POROSITY	.00000	.00000	1.00000		
HYDROCARBON SATURATION	.00000	.00000	.00000	1.00000	
GAS FRACTION	.00000	.00000	.00000	.00000	1.00000
GAS FWN VOLUME FACTOR	.00000	.00000	.00000	.00000	.00000

VARIABLE	
GAS FWN VOLUME FACTOR	1.00000

PETRIMES MODULE MPRO

NO. OF POOLS DISTRIBUTION AND RISKS
****

UAI CS4D0101
PLAY Herschel gas play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date FRI, JAN 26, 2001, 12:26 PM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE ON DB? > Y
OIL (O) OR GAS (G) ? > G

A) Risks

	GEOLOGICAL FACTOR		MARGINAL PROBABILITY
PLAY LEVEL	Overall Play Level Risk	=	1.00
PROSPECT LEVEL	Presence of Closure	(1)	.70
	Adequate Seal	(4)	.90
	Adequate Timing	(5)	.60
	Adequate Source	(6)	.80
	Adequate Maturation	(7)	.70
	Adequacy of Recovery	(9)	.80
	Overall Prospect Level Risk	=	.17
EXPLORATION RISK:		=	.17

B) No. of Prospects Distribution

Minimum = 10
Maximum = 50
Mean = 27.92
S.D. = 11.72

Frequency	No. of Prospects
99.00	10
95	12
90	13
80	16
75	18
60	22
50	25
40	30
25	38
20	40
10	45
5	48
1	50
0	50

C) No. of Pools Distribution

Minimum = 0
Maximum = 21
Mean = 4.73
S.D. = 2.80

Frequency	No. of Pools
97.40	0
95	1
90	1
80	2
75	3
60	4
50	4
40	5
25	6
20	7
10	8
5	10
1	12
0	21

Note: The no. of pools distribution is saved in the database with UDI= 6201GB4

PETRIMES MODULE PSRK

INDIVIDUAL POOL SIZES BY RANK
WHERE N IS A RANDOM VARIABLE
****

UAI CS4D0101
PLAY Herschel gas play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date FRI, JAN 26, 2001, 12:27 PM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE ON DB ? > Y
DO YOU WANT TO USE MPRO OUTPUT? > Y
MIN. AND MAX. POOL RANKS? > 1 5
DO YOU USE LOGNORNAL ASSUMPTION? > Y
DO YOU WANT TO USE PPSD OUTPUT? > Y

A) Basic Information

TYPE OF RESOURCE =Gas In-place
SYSTEM OF MEASUREMENT =S.I.
UNIT OF MEASUREMENT =M cu m (19)

B) Lognormal Pool Size Distribution

Summary Statistics	mu	= 7.9831	MEAN	= 4163.4		
	sig. sq=	.90202	S.D.	= 5038.5		
Upper Percentiles	99.99%	= 77.552	60.00%	= 2084.9	15.00%	= 7096.9
	99.00%	= 291.09	55.00%	= 2353.7	10.00%	= 8957.2
	95.00%	= 556.06	50.00%	= 2652.0	8.00%	= 10072.
	90.00%	= 785.19	45.00%	= 2980.2	6.00%	= 11611.
	85.00%	= 991.01	40.00%	= 3373.4	5.00%	= 12648.
	80.00%	= 1192.4	35.00%	= 3823.9	4.00%	= 13985.
	75.00%	= 1397.5	30.00%	= 4363.9	2.00%	= 18650.
	70.00%	= 1611.7	25.00%	= 5032.5	1.00%	= 24161.
	65.00%	= 1839.2	20.00%	= 5898.2	.01%	= 90689.

C) No. of Pools Distribution

Lower Support = 0
Upper Support = 21
Expectation = 4.73
Standard Deviation= 2.80

D) Pool Sizes By Rank

Pool Rank			Distribution			
1	MEAN	= 9269.1	S.D.	= 8208.5	P(N>=r)=	.97399
	99%	= 793.97	75%	= 4286.8	10%	= 18016.
	95%	= 1754.1	50%	= 7145.8	5%	= 23666.
	90%	= 2528.8	25%	= 11591.	1%	= 40378.
2	MEAN	= 4736.3	S.D.	= 3343.4	P(N>=r)=	.89343
	99%	= 464.17	75%	= 2412.7	10%	= 8885.1
	95%	= 967.94	50%	= 4000.9	5%	= 10976.
	90%	= 1398.3	25%	= 6184.8	1%	= 16296.
3	MEAN	= 3251.1	S.D.	= 2156.4	P(N>=r)=	.76446
	99%	= 353.87	75%	= 1692.0	10%	= 6037.4
	95%	= 698.41	50%	= 2806.1	5%	= 7317.5
	90%	= 990.25	25%	= 4295.9	1%	= 10372.
4	MEAN	= 2497.8	S.D.	= 1597.8	P(N>=r)=	.61665
	99%	= 299.30	75%	= 1325.3	10%	= 4598.1
	95%	= 568.84	50%	= 2176.4	5%	= 5522.6
	90%	= 791.83	25%	= 3307.3	1%	= 7653.6
5	MEAN	= 2033.5	S.D.	= 1264.1	P(N>=r)=	.47506
	99%	= 265.01	75%	= 1101.0	10%	= 3709.6
	95%	= 490.02	50%	= 1783.0	5%	= 4431.4
	90%	= 671.85	25%	= 2687.7	1%	= 6061.7

E) The mean of the potential = 18251.

PETRIMES MODULE PSUM

MONTE CARLO SUM SIMULATION
POOL SIZE DISTRIBUTION
****

UAI C56D0101
PLAY Herschel gas play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date FRI, JAN 26, 2001, 12:28 PM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE IN DATA BASE ?	>	Y
OIL (O) OR GAS (G) ?	>	G
BRITISH OR S.I. UNIT OF MEASUREMENT?	>	SI
RECOVERABLE RESOURCES?	>	N
DO YOU WANT TO USE MPRO OUTPUT?	>	Y
DO YOU ASSUME LOGNORMAL DISTRIBUTION?	>	Y
DO YOU WANT TO USE PPSD OUTPUT?	>	Y
DO YOU COMPUTE CONDITIONAL POTENTIAL?	>	N

A) Basic Information

TYPE OF RESOURCE =Gas In-place
SYSTEM OF MEASUREMENT =S.I.
UNIT OF MEASUREMENT =M cu m (19)

B) Lognormal Pool Size Distribution

Summary Statistics	mu	= 7.8831	MEAN	= 4163.4		
	sig. sq=	.90202	S.D.	= 5038.5		
Upper Percentiles	99.99%	= 77.552	60.00%	= 2084.9	15.00%	= 7096.9
	99.00%	= 291.09	55.00%	= 2353.7	10.00%	= 8957.2
	95.00%	= 556.06	50.00%	= 2652.0	8.00%	= 10072.
	90.00%	= 785.19	45.00%	= 2988.2	6.00%	= 11611.
	85.00%	= 991.01	40.00%	= 3373.4	5.00%	= 12648.
	80.00%	= 1192.4	35.00%	= 3823.9	4.00%	= 13985.
	75.00%	= 1397.5	30.00%	= 4363.9	2.00%	= 18650.
	70.00%	= 1611.7	25.00%	= 5032.5	1.00%	= 24161.
	65.00%	= 1839.2	20.00%	= 5898.2	.01%	= 90689.

C) NO. OF POOLS DISTRIBUTION

Lower Support	=	0
Upper Support	=	21
Expectation	=	4.72808
Standard Deviation	=	2.80484

D) Summary Statistics for 4000 Simulations

Play Resource: (B cu m)

Minimum = .0000000E+00 Maximum = 154.6978
Expectation = 19.58015 Standard Deviation = 15.77324

EMPERICAL DISTRIBUTION:

Greater than Percentage	Play Potential
100.00	.00000E+00
95.00	1.3186
90.00	3.2235
85.00	4.9252
80.00	6.3871
75.00	7.9166
70.00	9.4051
65.00	10.943
60.00	12.589
55.00	14.316
50.00	16.095
45.00	17.924
40.00	20.023
35.00	22.259
30.00	24.335
25.00	27.416
20.00	30.370
15.00	34.706
10.00	40.845
8.00	43.780
6.00	46.897
5.00	49.528
4.00	52.672
2.00	60.664
1.00	71.442
.01	144.59
.00	153.69

PETRIMES MODULE PPSD

PROSPECT OR POOL SIZE DISTRIBUTION USING LOGNORMAL OR MONTE CARLO SIMULATION

UAI	CS4K0101
PLAY	Yukon coastal plain gas play
Assessor	Peter Hannigan
Geologist	Peter Hannigan
Operator	Y
Remarks	Yukon North Slope Hydrocarbon Assessment Project
Run date	FRI, FEB 2, 2001, 12:00 PM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE ON DB?	> Y
OIL (O) OR GAS (G) ?	> G
BRITISH OR S.I. UNIT OF MEASUREMENT?	> SI
RECOVERABLE RESOURCES?	> N
DO YOU WANT TO HONOR A PERCENTILE?	> Y
UPPER PERCENTILE TO BE HONORED?	> .10000E-01

A) Basic Information

TYPE OF RESOURCE	=Gas In-place
SYSTEM OF MEASUREMENT	=S.I.
UNIT OF MEASUREMENT	=M cu m (19)

B) Input Constant(s)

No input constant.

C) Summary

Variable			Lognormal Approximation		
IN-PLACE	mu	= 7.9616	MEAN	= 4871.3	
POOL SIZE	sig. sq	= 1.0590	S.D.	= 6685.3	
(DERIVED)	99.99%	= 62.459	60.00%	= 2210.4	15.00% = 8334.8
	99.00%	= 261.82	55.00%	= 2520.8	10.00% = 10726.
	95.00%	= 527.93	50.00%	= 2868.8	8.00% = 12180.
	90.00%	= 767.26	45.00%	= 3264.8	6.00% = 14209.
	85.00%	= 987.40	40.00%	= 3723.2	5.00% = 15589.
	80.00%	= 1206.6	35.00%	= 4264.8	4.00% = 17382.
	75.00%	= 1433.0	30.00%	= 4921.0	2.00% = 23744.
	70.00%	= 1672.4	25.00%	= 5742.9	1.00% = 31433.
	65.00%	= 1929.7	20.00%	= 6820.7	.01% = .13176E+06

D) Pool Size Equation

NO. OF GEO-VARIABLES= 6

POOL SIZE = 1.00000
* AREA OF CLOSURE * FORMATION THICKNESS * POROSITY * TRAP FILL
* HYDROCARBON SATURATION

GAS FVN VOLUME FACTOR

GEOLOGICAL VARIABLES

A) Variables Used

VARIABLE	UNIT OF MEASUREMENT SYMBOL	O/G	VERSION	TYPE	NO. OF POINTS USED	UAI
AREA OF CLOSURE	sq km	Gas	1	2	4	C54K0101
FORMATION THICKNESS	m	Gas	1	2	4	C54K0101
POROSITY	dec fr	Gas	1	2	4	C54K0101
TRAP FILL	dec fr	Gas	1	2	4	C54K0101
HYDROCARBON SATURATION	dec fr	Gas	1	2	4	C54K0101
GAS FVN VOLUME FACTOR		Gas	1	2	4	C54K0101

B) Summary

Variable — Lognormal Approximation

AREA OF CLOSURE	mu = 2.1972	MEAN = 12.505			
	sig. sq= .65782	S.D. = 12.063			
	99.99% = .44082	60.00% = 7.3283	15.00% = 20.860		
	99.00% = 1.3640	55.00% = 8.1279	10.00% = 25.448		
	95.00% = 2.3706	50.00% = 9.0000	8.00% = 28.130		
	90.00% = 3.1829	45.00% = 9.9656	6.00% = 31.761		
	85.00% = 3.8830	40.00% = 11.053	5.00% = 34.168		
	80.00% = 4.5477	35.00% = 12.302	4.00% = 37.231		
	75.00% = 5.2079	30.00% = 13.771	2.00% = 47.605		
	70.00% = 5.8820	25.00% = 15.553	1.00% = 59.385		
	65.00% = 6.5844	20.00% = 17.811	.01% = 183.75		
FORMATION THICKNESS	mu = 4.3175	MEAN = 82.958			
	sig. sq= .20151	S.D. = 39.194			
	99.99% = 14.126	60.00% = 66.930	15.00% = 119.43		
	99.00% = 26.395	55.00% = 70.886	10.00% = 133.32		
	95.00% = 35.842	50.00% = 75.000	8.00% = 140.92		
	90.00% = 42.191	45.00% = 79.352	6.00% = 150.72		
	85.00% = 47.098	40.00% = 84.034	5.00% = 156.94		
	80.00% = 51.402	35.00% = 89.162	4.00% = 164.58		
	75.00% = 55.407	30.00% = 94.906	2.00% = 188.56		
	70.00% = 59.269	25.00% = 101.52	1.00% = 213.11		
	65.00% = 63.087	20.00% = 109.43	.01% = 398.20		
POROSITY	mu = -2.3026	MEAN = .10226			
	sig. sq= .44688E-01	S.D. = .21861E-01			
	99.99% = .45558E-01	60.00% = .94785E-01	15.00% = .12450		
	99.00% = .61154E-01	55.00% = .97379E-01	10.00% = .13112		
	95.00% = .70630E-01	50.00% = .10000	8.00% = .13458		
	90.00% = .76268E-01	45.00% = .10269	6.00% = .13891		
	85.00% = .80324E-01	40.00% = .10550	5.00% = .14150		
	80.00% = .83701E-01	35.00% = .10849	4.00% = .14479		
	75.00% = .86711E-01	30.00% = .11172	2.00% = .15437		
	70.00% = .89507E-01	25.00% = .11532	1.00% = .16352		
	65.00% = .92177E-01	20.00% = .11947	.01% = .21950		
TRAP FILL	mu = -1.3863	MEAN = .26798			
	sig. sq= .13819	S.D. = .10312			
	99.99% = .62738E-01	60.00% = .22753	15.00% = .36751		
	99.00% = .10529	55.00% = .23859	10.00% = .40257		
	95.00% = .13564	50.00% = .25000	8.00% = .42148		
	90.00% = .15525	45.00% = .26196	6.00% = .44560		
	85.00% = .17007	40.00% = .27469	5.00% = .46078		
	80.00% = .18204	35.00% = .28850	4.00% = .47926		
	75.00% = .19456	30.00% = .30381	2.00% = .53642		
	70.00% = .20572	25.00% = .32124	1.00% = .59362		
	65.00% = .21664	20.00% = .34183	.01% = .99620		
HYDROCARBON SATURATION	mu = -.16252	MEAN = .85038			
	sig. sq= .88600E-03	S.D. = .25318E-01			
	99.99% = .76093	60.00% = .84361	15.00% = .87663		
	99.00% = .79313	55.00% = .84683	10.00% = .88305		
	95.00% = .80939	50.00% = .85000	8.00% = .88630		
	90.00% = .81819	45.00% = .85319	6.00% = .89026		
	85.00% = .82418	40.00% = .85643	5.00% = .89265		
	80.00% = .82897	35.00% = .85981	4.00% = .89547		
	75.00% = .83311	30.00% = .86337	2.00% = .90358		
	70.00% = .83684	25.00% = .86724	1.00% = .91094		
	65.00% = .84031	20.00% = .87156	.01% = .94950		
GAS FVN VOLUME FACTOR	mu = -5.2983	MEAN = .50399E-02			
	sig. sq= .15887E-01	S.D. = .63777E-03			
	99.99% = .31289E-02	60.00% = .48429E-02	15.00% = .56978E-02		
	99.00% = .37293E-02	55.00% = .49214E-02	10.00% = .58765E-02		
	95.00% = .40638E-02	50.00% = .50000E-02	8.00% = .59687E-02		
	90.00% = .42542E-02	45.00% = .50798E-02	6.00% = .60824E-02		
	85.00% = .43877E-02	40.00% = .51622E-02	5.00% = .61519E-02		
	80.00% = .44968E-02	35.00% = .52488E-02	4.00% = .62345E-02		
	75.00% = .45925E-02	30.00% = .53416E-02	2.00% = .64773E-02		
	70.00% = .46802E-02	25.00% = .54437E-02	1.00% = .67037E-02		
	65.00% = .47630E-02	20.00% = .55596E-02	.01% = .79903E-02		

C) Covariance Matrix

VARIABLE					
AREA OF CLOSURE	.65782				
FORMATION THICKNESS	.00000	.20151			
POROSITY	.00000	.00000	.04469		
TRAP FILL	.00000	.00000	.00000	.13819	
HYDROCARBON SATURATION	.00000	.00000	.00000	.00000	.00089
GAS FVH VOLUME FACTOR	.00000	.00000	.00000	.00000	.00000

VARIABLE	
GAS FVH VOLUME FACTOR	.01589

D) Correlation Matrix

VARIABLE					
AREA OF CLOSURE	1.00000				
FORMATION THICKNESS	.00000	1.00000			
POROSITY	.00000	.00000	1.00000		
TRAP FILL	.00000	.00000	.00000	1.00000	
HYDROCARBON SATURATION	.00000	.00000	.00000	.00000	1.00000
GAS FVH VOLUME FACTOR	.00000	.00000	.00000	.00000	.00000

VARIABLE	
GAS FVH VOLUME FACTOR	1.00000

PETRIMES MODULE MPRO

NO. OF POOLS DISTRIBUTION AND RISKS

UAI C54K0101
PLAY Yukon coastal plain gas play
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date FRI, FEB 2, 2001, 11:47 AM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE ON DB? > Y
OIL (O) OR GAS (G) ? > G

A) Risks

	GEOLOGICAL FACTOR		MARGINAL PROBABILITY
PLAY LEVEL	Overall Play Level Risk	=	1.00
PROSPECT LEVEL	Presence of Closure	(1)	.70
	Presence of Porosity	(3)	.65
	Adequate Seal	(4)	.80
	Adequate Timing	(5)	.60
	Adequate Source	(6)	.80
	Adequate Preservation	(8)	.60
	Overall Prospect Level Risk	=	.10
EXPLORATION RISK:		=	.10

B) No. of Prospects Distribution

Minimum = 5
Maximum = 40
Mean = 24.15
S.D. = 10.25

Frequency	No. of Prospects
99.00	5
95	7
90	9
80	13
75	15
60	21
50	25
40	28
25	33
20	34
10	37
5	39
1	40
0	40

C) No. of Pools Distribution

Minimum = 0
Maximum = 14
Mean = 2.53
S.D. = 1.85

Frequency	No. of Pools
87.27	0
80	1
75	1
60	2
50	2
40	3
25	4
20	4
10	5
5	6
1	8
0	14

Note: The no. of pools distribution is saved in the database with UDI= 6201GB4

PETRIMES MODULE PSRK

```
INDIVIDUAL POOL SIZES BY RANK
WHERE N IS A RANDOM VARIABLE
*****************************
UAI         C54K0101
PLAY        Yukon coastal plain gas play
Assessor    Peter Hannigan
Geologist   Peter Hannigan
Remarks     Yukon North Slope Hydrocarbon Assessment Project
Run date    FRI, FEB  2, 2001, 12:02 PM

USER SUPPLIED PARAMETERS
------------------------
   DO YOU WANT TO STORE ON DB ?          > Y
   DO YOU WANT TO USE MPRO OUTPUT?       > Y
   MIN. AND MAX. POOL RANKS?             >      1     3
   DO YOU USE LOGNORNAL ASSUMPTION?      > Y
   DO YOU WANT TO USE PPSD OUTPUT?       > Y

A) Basic Information
   -----------------
   TYPE OF RESOURCE      =Gas In-place
   SYSTEM OF MEASUREMENT =S.I.
   UNIT OF MEASUREMENT   =M cu m  (19)

B) Lognormal Pool Size Distribution
   --------------------------------
   Summary      mu     = 7.9616     MEAN   = 4871.3
   Statistics   sig. sq= 1.0590     S.D.   = 6685.3
   Upper        99.99% = 62.459     60.00% = 2210.4     15.00% = 8334.8
   Percentiles  99.00% = 261.82     55.00% = 2520.8     10.00% = 10726.
                95.00% = 527.93     50.00% = 2868.8      8.00% = 12180.
                90.00% = 767.26     45.00% = 3264.8      6.00% = 14209.
                85.00% = 987.40     40.00% = 3723.2      5.00% = 15589.
                80.00% = 1206.6     35.00% = 4264.8      4.00% = 17382.
                75.00% = 1433.0     30.00% = 4921.0      2.00% = 23744.
                70.00% = 1672.4     25.00% = 5742.9      1.00% = 31433.
                65.00% = 1929.7     20.00% = 6820.7       .01% = .13176E+06

C) No. of Pools Distribution
   -------------------------
   Lower Support      =  0
   Upper Support      =  14
   Expectation        =   2.53
   Standard Deviation=   1.85

D) Pool Sizes By Rank
   ------------------
   Pool Rank                        Distribution
        1     MEAN   = 8704.3      S.D.   = 9663.2     P(N>=r)= .87266
              99%    = 475.24      75%    = 3186.3     10%    = 18198.
              95%    = 1092.3      50%    = 5998.4      5%    = 25009.
              90%    = 1670.0      25%    = 10790.      1%    = 46181.
        2     MEAN   = 3994.3      S.D.   = 3404.2     P(N>=r)= .66507
              99%    = 311.26      75%    = 1717.3     10%    = 8047.4
              95%    = 647.01      50%    = 3093.3      5%    = 10348.
              90%    = 946.04      25%    = 5211.9      1%    = 16478.
        3     MEAN   = 2601.7      S.D.   = 2023.4     P(N>=r)= .45238
              99%    = 245.65      75%    = 1194.3     10%    = 5132.0
              95%    = 483.20      50%    = 2085.8      5%    = 6457.0
              90%    = 685.68      25%    = 3422.8      1%    = 9775.8

E) The mean of the potential =  11429.
```

PETRIMES MODULE PSUM

```
MONTE CARLO SUM SIMULATION
POOL SIZE DISTRIBUTION
**************************

UAI         C54K0101
PLAY        Yukon coastal plain gas play
Assessor    Peter Hannigan
Geologist   Peter Hannigan
Remarks     Yukon North Slope Hydrocarbon Assessment Project
Run date    FRI, FEB  2, 2001, 12:03 PM

USER SUPPLIED PARAMETERS
------------------------
   DO YOU WANT TO STORE IN DATA BASE ?   >   Y
   OIL (O) OR GAS (G) ?                  >   G
   BRITISH OR S.I. UNIT OF MEASUREMENT?  >   SI
   RECOVERABLE RESOURCES?                >   N
   DO YOU WANT TO USE MPRO OUTPUT?       >   Y
   DO YOU ASSUME LOGNORMAL DISTRIBUTION? >   Y
   DO YOU WANT TO USE PPSD OUTPUT?       >   Y
   DO YOU COMPUTE CONDITIONAL POTENTIAL? >   N

A) Basic Information
   -----------------
   TYPE OF RESOURCE      =Gas In-place
   SYSTEM OF MEASUREMENT =S.I.
   UNIT OF MEASUREMENT   =M cu m  (19)

B) Lognormal Pool Size Distribution
   --------------------------------
   Summary      mu     = 7.9616     MEAN   = 4871.3
   Statistics   sig. sq= 1.0590     S.D.   = 6685.3

   Upper        99.99% = 62.459     60.00% = 2210.4     15.00% = 8334.8
   Percentiles  99.00% = 261.82     55.00% = 2520.8     10.00% = 10726.
                95.00% = 527.93     50.00% = 2868.8      8.00% = 12180.
                90.00% = 767.26     45.00% = 3264.8      6.00% = 14209.
                85.00% = 987.40     40.00% = 3723.2      5.00% = 15589.
                80.00% = 1206.6     35.00% = 4264.8      4.00% = 17382.
                75.00% = 1433.0     30.00% = 4921.0      2.00% = 23744.
                70.00% = 1672.4     25.00% = 5742.9      1.00% = 31433.
                65.00% = 1929.7     20.00% = 6820.7       .01% = .13176E+06

C) NO. OF POOLS DISTRIBUTION
   -------------------------
   Lower Support      =  0
   Upper Support      =  14
   Expectation        =   2.53117
   Standard Deviation=   1.84960
```

D) *Summary Statistics for 4000 Simulations

Play Resource: (B cu m)

Minimum = .0000000E+00 Maximum = 157.3106
Expectation = 12.44228 Standard Deviation= 13.61104

EMPERICAL DISTRIBUTION:

Greater than Percentage	Play Potential
100.00	.00000E+00
85.00	.76783
80.00	1.7893
75.00	2.8329
70.00	3.8510
65.00	4.8858
60.00	6.0649
55.00	7.2271
50.00	8.4634
45.00	9.8860
40.00	11.438
35.00	13.128
30.00	15.289
25.00	17.542
20.00	20.512
15.00	24.260
10.00	29.722
8.00	32.826
6.00	36.776
5.00	38.204
4.00	40.867
2.00	50.116
1.00	59.716
.01	151.39
.00	156.72

PETRIMES MODULE PSUM

MONTE CARLO SUM SIMULATION
POOL SIZE DISTRIBUTION
****

UAI C5410101
PLAY All oil plays
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date MON, JAN 29, 2001, 2:16 PM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE IN DATA BASE ? > Y
OIL (O) OR GAS (G) ? > O
BRITISH OR S.I. UNIT OF MEASUREMENT? > SI
RECOVERABLE RESOURCES? > N
DO YOU COMPUTE CONDITIONAL POTENTIAL? > N

A) Basic Information

TYPE OF RESOURCE =Oil In-place
SYSTEM OF MEASUREMENT =S.I.
UNIT OF MEASUREMENT =M cu m (19)

B) PLAY POTENTIAL DISTRIBUTION

Summary Statistics: MEAN = 3.7985 S.D. = 3.6877 M cu m

Upper Percentiles	100.00% = .00000E+00	55.00% = 2.5113	10.00% = 8.1451
	95.00% = .24337	50.00% = 2.8268	8.00% = 8.8912
	90.00% = .55668	45.00% = 3.1302	6.00% = 9.7734
	85.00% = .84104	40.00% = 3.5563	5.00% = 10.497
	80.00% = 1.0941	35.00% = 3.9955	4.00% = 11.326
	75.00% = 1.3875	30.00% = 4.4510	2.00% = 14.251
	70.00% = 1.6443	25.00% = 5.1129	1.00% = 16.738
	65.00% = 1.9165	20.00% = 5.7703	.01% = 46.342
	60.00% = 2.1850	15.00% = 6.6875	.00% = 47.536

Summary Statistics: MEAN = 42.406 S.D. = 34.551 M cu m

Upper Percentiles	100.00% = .00000E+00	55.00% = 30.753	10.00% = 88.821
	95.00% = 2.7805	50.00% = 34.605	8.00% = 95.161
	90.00% = 6.8992	45.00% = 38.698	6.00% = 101.91
	85.00% = 10.490	40.00% = 43.242	5.00% = 107.90
	80.00% = 13.667	35.00% = 48.156	4.00% = 114.67
	75.00% = 16.913	30.00% = 52.590	2.00% = 132.80
	70.00% = 20.214	25.00% = 59.358	1.00% = 156.25
	65.00% = 23.473	20.00% = 65.791	.01% = 324.33
	60.00% = 27.095	15.00% = 75.218	.00% = 346.56

C) NO. OF PLAYS DISTRIBUTION

Lower Support = 2
Upper Support = 2
Expectation = 2.00000
Standard Deviation= .00000

D) Summary Statistics for 4000 Simulations

Basin Resource: (M cu m)

Minimum = .0000000E+00 Maximum = 328.2340
Expectation = 47.60648 Standard Deviation= 37.01921

EMPERICAL DISTRIBUTION:

Greater than Percentage	Basin Potential
100.00	.00000E+00
99.00	1.3516
95.00	6.6909
90.00	11.123
85.00	14.978
80.00	18.259
75.00	21.697
70.00	25.456
65.00	28.812
60.00	32.447
55.00	35.995
50.00	39.767
45.00	43.931
40.00	48.463
35.00	52.726
30.00	57.466
25.00	63.285
20.00	71.387
15.00	80.399
10.00	91.935
8.00	99.354
6.00	105.94
5.00	110.98
4.00	118.50
2.00	137.32
1.00	150.43
.01	326.42
.00	328.05

PETRIMES MODULE PSUM

MONTE CARLO SUM SIMULATION
POOL SIZE DISTRIBUTION
****

UAI C5400101
PLAY All gas plays
Assessor Peter Hannigan
Geologist Peter Hannigan
Remarks Yukon North Slope Hydrocarbon Assessment Project
Run date FRI, FEB 2, 2001, 12:09 PM

USER SUPPLIED PARAMETERS

DO YOU WANT TO STORE IN DATA BASE ? > Y
OIL (O) OR GAS (G) ? > G
BRITISH OR S.I. UNIT OF MEASUREMENT? > SI
RECOVERABLE RESOURCES? > N
DO YOU COMPUTE CONDITIONAL POTENTIAL? > N

A) Basic Information

TYPE OF RESOURCE =Gas In-place
SYSTEM OF MEASUREMENT =S.I.
UNIT OF MEASUREMENT =M cu m (19)

B) PLAY POTENTIAL DISTRIBUTION

Summary Statistics	MEAN = 2.4325 B cu m		S.D. = 1.7427			
Upper Percentiles	100.00% = .00000E+00		55.00% = 1.8698		8.00% = 4.9254	
	99.00% = .18142		50.00% = 2.0402		6.00% = 5.3993	
	95.00% = .48941		45.00% = 2.2209		5.00% = 5.6884	
	90.00% = .72355		40.00% = 2.4170		4.00% = 6.0976	
	85.00% = .92550		35.00% = 2.6301		2.00% = 7.0300	
	80.00% = 1.0815		30.00% = 2.8758		1.00% = 8.2659	
	75.00% = 1.2620		25.00% = 3.1710		.01% = 10.688	
	70.00% = 1.4062		20.00% = 3.5065		.00% = 20.140	
	65.00% = 1.5508		15.00% = 3.9387			
	60.00% = 1.7089		10.00% = 4.5919			
Summary Statistics	MEAN = 7.5234 B cu m		S.D. = 4.6634			
Upper Percentiles	100.00% = .15446		55.00% = 6.0653		8.00% = 14.470	
	99.00% = .87468		50.00% = 6.6614		6.00% = 15.512	
	95.00% = 1.6994		45.00% = 7.2674		5.00% = 16.536	
	90.00% = 2.3555		40.00% = 7.8857		4.00% = 17.344	
	85.00% = 2.9483		35.00% = 8.5103		2.00% = 19.535	
	80.00% = 3.4788		30.00% = 9.2216		1.00% = 21.752	
	75.00% = 3.9972		25.00% = 10.064		.01% = 33.164	
	70.00% = 4.5850		20.00% = 11.098		.00% = 33.429	
	65.00% = 5.0365		15.00% = 12.215			
	60.00% = 5.5197		10.00% = 13.715			
Summary Statistics	MEAN = 19.580 B cu m		S.D. = 15.773			
Upper Percentiles	100.00% = .00000E+00		55.00% = 14.314		10.00% = 40.845	
	95.00% = 1.3186		50.00% = 16.095		8.00% = 43.780	
	90.00% = 3.2235		45.00% = 17.924		6.00% = 46.897	
	85.00% = 4.9252		40.00% = 20.023		5.00% = 49.528	

```
      80.00% = 6.3871          35.00% = 22.259          4.00% = 52.672
      75.00% = 7.9166          30.00% = 24.335          2.00% = 60.664
      70.00% = 9.4051          25.00% = 27.416          1.00% = 71.442
      65.00% = 10.943          20.00% = 30.370           .01% = 144.59
      60.00% = 12.589          15.00% = 34.706           .00% = 153.69

   Summary      MEAN   = 12.442     S.D.   = 13.611
   Statistics   B cu m

   Upper      100.00% = .00000E+00  45.00% = 9.8860      6.00% = 36.776
   Percentiles 85.00% = .76783      40.00% = 11.438      5.00% = 38.204
               80.00% = 1.7893      35.00% = 13.128      4.00% = 40.867
               75.00% = 2.8329      30.00% = 15.289      2.00% = 50.116
               70.00% = 3.8510      25.00% = 17.542      1.00% = 59.716
               65.00% = 4.8850      20.00% = 20.512       .01% = 151.39
               60.00% = 6.0649      15.00% = 24.260       .00% = 156.72
               55.00% = 7.2271      10.00% = 29.722
               50.00% = 8.4634       8.00% = 32.826

C) NO. OF PLAYS DISTRIBUTION
   ---------------------------
   Lower Support      =  4
   Upper Support      =  4
   Expectation        =  4.00000
   Standard Deviation=   .00000

D) Summary Statistics for 4000 Simulations
   -----------------------------------------

   Basin Resource:  ( B cu m )
   ---------------------------
     Minimum       = 3.050182   Maximum                = 194.6945
     Expectation = 42.47473     Standard Deviation= 22.73458

   EMPERICAL DISTRIBUTION:
   -----------------------
     Greater than   Basin
     Percentage     Potential
     ----------     ---------
        100.00        3.0302
         99.00        8.3587
         95.00       14.470
         90.00       18.727
         85.00       21.995
         80.00       24.276
         75.00       26.703
         70.00       29.099
         65.00       31.370
         60.00       33.781
         55.00       36.138
         50.00       38.565
         45.00       41.272
         40.00       44.074
         35.00       46.562
         30.00       49.632
         25.00       53.034
         20.00       57.403
         15.00       62.927
         10.00       69.836
          8.00       73.161
          6.00       77.939
          5.00       81.829
          4.00       85.455
          2.00      102.55
          1.00      123.30
              .01   194.01
              .00   194.63
```